



AMERICAN JOURNAL OF PHARMTECH RESEARCH

Journal home page: <http://www.ajptr.com/>

An evaluation of tuberculosis cases: A retrospective study

Bhimaray Krishnagoudar*¹, Sandeep Anandmurthy¹, Katti Ravi Venkappa¹,
Mahadevamma L¹, Shaik Shafiya Begum¹

*1. Department of Pharmacy Practice, Sri Adichunchanagiri College of Pharmacy, B.G.Nagara,
Karnataka -571448. India.*

ABSTRACT

Tuberculosis is a contagious infection caused by air borne bacteria Mycobacterium tuberculosis. Tuberculosis is a growing health problem in the developing world. India accounts for one-fifth of the global TB incident cases, each year nearly two million people in India develop TB¹. This was a retrospective record based survey carried at AH & RC (Adichunchanagiri Hospital & Research Centre) Tertiary care teaching hospital, B.G Nagara. Twelve month data of all TB patients i.e. from Sep-09 to Aug-10 that were diagnosed in the Directly Observed Treatment Short Course Centre was taken, which included 120 diagnosed patients. Out of 120 patients diagnosed, the male to female ratio was 2.5:1, and 29 were from age group of 40 to 49 years. Pulmonary TB cases were more i.e. 85 (75.83%) when compared to extra pulmonary i.e. 35 (29.17%), new smear positive cases were 62 (51.67%) and new smear negative cases were 58 (48.33%). Total 67 (55.83%) patients were categorized in CAT-I, 25 (20.83%) patients in CAT-II and 28 (23.33%) in CAT-III. The Treatment Completion Rate (TCR) and rate of cure was not known since all patients were transferred to their nearest peripheral RNTCP/DOTS Centers, and those centers failed to provide proper feedback. So, for transferred TB cases a better system of follow up should be done in order to know about the TCR and rate of cure.

Key Words: RNTCP/DOTS centre, Category, Tuberculosis, TCR (Treatment Completion Rate).

*Corresponding Author Email: bsk.lkp@gmail.com

Received 8 April 2012, Accepted 28 April 2012

Please cite this article in press as: Krishnagoudar B *et al.*, An evaluation of tuberculosis cases: A retrospective study. American Journal of PharmTech Research 2012.

INTRODUCTION

Worldwide, Mycobacterium tuberculosis remains the leading infective cause of mortality and morbidity¹. Each year about 8 million people worldwide develop Tuberculosis(TB). Of these, 95% occur in developing countries. India accounts for about 20%of the global burden of TB. Nearly 2.2 million people develop TB and about 3.5 lakh die every year².

Tuberculosis is a social disease with medical implications. It has always occurred disproportionately among disadvantaged populations such as the homeless, malnourished, and over-crowded. Within the past decade it also has become clear that the spread of HIV infection and the immigration of persons from areas of high incidence have resulted in increased numbers of TB cases³.

The National Tuberculosis Control Programme (NTP) was implemented in 1962. However, when reviewed in 1992, after three decades of implementation, the NTP was shown to have made no epidemiological impact, mainly due to poor case finding and low treatment completion rates⁵. As a result, the Government of India (GoI) in 1993 developed the Revised National Tuberculosis Control Programme (RNTCP) based on the internationally recommended Directly Observed Treatment – Short course (DOTS) strategy⁵.

Hospital based studies show that most of the patients admitted to the hospital are middle aged males. Pulmonary TB is the most common form of TB (more than 85% of all TB cases) while Extra-pulmonary TB can affect almost any organ in the body. The prevalence of TB infection in India is 30%. As per statement of Director General of WHO, DOTS strategy was most important breakthrough of the decade in terms of lives saved. The WHO. recommended DOTS strategy (Directly Observed Treatment Short Course) was covered in the entire India by 24th March 2006⁶.The past decade has seen a rapid expansion of DOTS centers throughout India, under the guidance of the RNTCP¹. Continued success of the programme, however, remains a daunting task as there has been very little co-ordination between the RNTCP, faculties of medical schools and health providers in the private sector. In medical schools, training on TB and HIV/AIDS does not focus on the public health aspects relating to prevention and control of these diseases⁴.

Pulmonary Tuberculosis (PTB):

The essential basis of diagnosis of pulmonary TB was three sputum smear examinations for M. tuberculosis (two spot and one early morning sample) according to the RNTCP guidelines.

Extra pulmonary Tuberculosis (EPTB):

Was diagnosed by the following criteria: (i) constitutional and organ specific clinical features depending upon the site of TB; (ii) radiographic features suggestive of TB; (iii) microbiological and /or histopathological diagnosis depending on the site (not an essential criteria for mediastinal lymph nodes and pleural effusion) and (iv) a satisfactory response to anti-TB treatment.

Categorization of Patients:

Patients were categorized into various treatment categories as per the RNTCP guidelines. Briefly, new sputum smear positive and seriously ill sputum smear negative pulmonary TB and extra pulmonary TB patients (Category I) were treated during intensive phase with four drugs: isoniazid (H), rifampicin (R), pyrazinamide (Z) and ethambutol (E) for two months followed by continuation phase of four months of two drugs, H and R. New sputum smear negative pulmonary TB and not seriously ill extra pulmonary TB patients (category III) were treated during the intensive phase with three drugs, H,R,Z for two months followed by continuation phase of four months of two drugs, H and R. Previously treated patients (category II) were treated during the intensive phase with five drugs H,R,Z,E and streptomycin (S) of initial two months and one month with four drugs as previously mentioned except S, followed by the continuation phase of five months with H, R and E. Smear positive patients whose smear was positive at the end of intensive phase received another month of intensive phase treatment. During the intensive phase every dose was given thrice weekly on alternate days under direct observation. Medications for continuation phase were packaged into weekly blister pack, the first dose of which was given under direct observation. The empty blister pack was returned the following weeks as evidence that the patient had taken the medicines⁴.

This study was carried out with the objectives -to estimate the number of New Smear Positive (NSP) case detected during diagnosis, -to study the age & sex distribution of TB patient diagnosed, -to study the variants of TB i.e. New Smear Positive, New Smear Negative, and Extra pulmonary TB, -categorization of all TB cases diagnosed at DOTS center.

MATERIAL AND METHODS

This was a retrospective record based survey carried out in the tertiary care teaching hospital of the rural area of south India. Adichunchanagiri is a 750 bedded tertiary care teaching hospital affiliated to Adichunchanagiri institute of medical sciences. RNTCP being a part of AH & RC, the patients diagnosed with TB are referred to the DOTS clinic where they are registered and treated according to RNTCP guidelines.

Data collection: Prior to collection of data from the RNTCP centre, permission was obtained from the medical superintendent of the hospital by explaining him the nature of the study. Then the record of 120 TB patients registered in the DOTS clinic from September 2009 to August 2010, were included in the study. Patients' registration numbers were used to obtain corresponding files from the medical records department. From each medical case file, the patient's history, physical findings, chest radiographs and reports of laboratory investigations were reviewed to obtain the necessary information about diagnosis of TB. The patients were divided into EPTB and PTB groups. The two groups were compared in respect to age, sex, and into case categories (new case or previously treated case). The new smear positive and smear negative cases were recorded in the suitable tabular form.

RESULTS AND DISCUSSION

Out of 120 patients diagnosed, the male to female ratio was 2.5:1, and the prevalence of TB was found more in the age group of 40 to 49 years, the total number of patients in this age group were 29, and the least was found to be in 10 to 19 years of age. Pulmonary TB cases were more i.e. 85 (75.83%) when compared to extra pulmonary TB cases i.e. 35 (29.17%), new smear positive cases were 62 (51.67%) and new smear negative cases were 58 (48.33%). Total 67 (55.83%) patients were categorized under Category-I, 25 (20.83%) patients in Category-II and 28 (23.33%) in Category-III.

Table 1 show that 71.66% males were diagnosed for TB compared to 28.33% of females. TB patients diagnosed in the age group of 40-49 years were more i.e. 29 (24.16%) out of this 20 were male and 09 females respectively. The next high prevalence rate was seen under the age group of 30-39 years i.e. 25 (20.83%), followed by 50-59 (19), 60-69 (16), 20-29 (13), 0-9 (8), 70-79 (6) years of age with percentages of 15.83%, 13.33%, 10.83%, 06.66%, and 05.00% respectively. There was increase in smear positivity due to the sensitization workshop organized for staff, postgraduates, nurses & interns. Out of 120 cases diagnosed, Category –I patients were 67 (55.83%) which include new smear positive, new smear negative and extra pulmonary cases. In 120 patients the age group between 30 to 70 was predominant with tuberculosis.

Table 2 shows that there was an drastic increase in diagnosis of new smear positive cases in the DOTS centre. Average diagnosis of new smear positive was 12.92%, with the male population being 50 (41.67%) and females 12 (10%). During the period of second quarter i.e. December-2009 to February-2010, 19 (30.65%) smear positive cases out of the total 62 new smear positive

cases were diagnosed, which was high when compared to other time periods. A very least i.e. 10 (08.33%) cases were identified during the third quarter of the study March-2010 to May-2010.

Table-1: Age and Sex distribution of tuberculosis patient.

Age (years)	Total patient diagnosed at RNTCP of AH & RC	
	Male	Female
0 - 9	05	03
10 - 19	02	02
20 - 29	10	03
30 - 39	17	08
40 - 49	20	09
50 - 59	13	06
60 - 69	14	02
70 - 79	05	01
	86(71.66%)	34(28.33%)
Grand total	120	

Table-2: Quarterly report of sputum evaluation (Sep-09 to Aug -10)

Quarter	New smear positive(NSP)		New smear negative (NSN)		Total
	Male	Female	Male	Female	
	Sep-09 to Nov-09	11	04	12	
Dec -09 to Feb -10	17	02	12	06	37
Mar-10 to May-10	08	02	04	09	23
June-10 to Aug -10	14	04	08	06	32
Grand total	50	12	36	22	120

The physician confirmed the diagnosis of tuberculosis in patients with two positive sputum smear specimens for acid-fast bacilli (AFB). They were further categorized as new (category I or category III) or old cases (category II) based on their treatment history as per the RNTCP criteria. If all the three sputum smear samples were negative for AFB and there was no response to one to two weeks of antibiotics, chest radiograph was taken and if consistent with TB, the patient was treated for smear negative TB. If only one of these samples was positive, chest radiograph was taken and patient was evaluated for pulmonary tuberculosis⁴.

Table 3 show about the type of TB diagnosed. The most cases found out were the pulmonary TB cases with 85 (70.83%) and extra pulmonary cases being 35 (29.17%). The pulmonary TB cases with age group of 30-39 were 20 (23.5%) and more compared to extra-pulmonary TB with the age group of 40-49 years with 12 (34.29%) cases. The pulmonary TB cases in the age group of 10-19 was least with 3 (02.5%).

Table 3: Pulmonary and Extrapulmonary TB

Age (years)	Total number of diagnosed TB		Pulmonary TB		Extra pulmonary TB	
	Male	Female	Male	Female	Male	Female
0 – 9	05	03	03	02	02	01
10 – 19	02	02	02	01	00	01
20 – 29	10	03	05	03	05	00
30 – 39	17	08	14	06	03	02
40 – 49	20	09	14	03	06	06
50 – 59	13	06	13	02	00	04
60 – 69	14	02	11	02	03	00
70 – 79	05	01	04	00	01	01
Total	86	34	66	19	20	15

All diagnosed TB patients were divided in three categories for therapeutic purposes adopted under RNTCP. Category 1 includes new sputum smear positive cases, seriously ill sputum smear negative cases and seriously ill extra pulmonary cases. Category 2 includes sputum positive relapse, sputum positive failure and treatment positive treatment after default cases. Category 3 includes sputum negative and extra pulmonary not seriously ill cases⁶.

Table 4 show that 67(55.83%) patients were in Category I, 25(20.83%) patients were in Category II and 28(23.33%) patients were in Category III. AH&RC being a tertiary care centre, more number of seriously ill patients were placed in Category I.

Table 4: Category wise distribution of TB patients

Categorization	CAT-I		CAT-II		CAT-III		Total
	Male	Female	Male	Female	Male	Female	
Total diagnosed	53	14	20	05	13	15	120

The patients whose smear was examined for TB showed the results with both 'smear positive' and 'smear negative'. But, for 'smear negative' patients empirical treatment with anti-TB drugs was given. As the diagnosed patients did not stay in the AH & RC centre until the treatment completion, such patients were put on DOTS and referred to their nearest peripheral RNTCP centre for the treatment completion. So, the inpatients data in the AH & RC centre was not available. The feedback of patients referred to the peripheral centers was not available due to lack of proper communication from the peripheral centers which hindered the treatment completion rate. There is a variation in the cases diagnosed at the centre due to poor health literacy among the population where most of the people are illiterate, even some of the patients gave wrong address or some have shifted from the previous residing area.

CONCLUSION

RNTCP/DOTS regimens were found to be effective to diagnose and treat even the complicated cases of TB. The TCR and rate of cure was not known since all patients were transferred to their nearest peripheral DOTS Centers and those centers failed to provide proper feedback. So, for transferred cases a better system of follow up should be done in order to know about the TCR and rate of cure.

ACKNOWLEDGEMENT

We thank Lokesh AC, TB health visitor, Gopal DP, lab technician who helped in providing tuberculosis case records. Thanks due to Dr. M Shivanna, Medical superintendent; also we thank Dr. B Ramesh Principal SAC College of Pharmacy.

REFERENCES

1. Sreeramreddy CT, Panduru KV, Verma SC. Comparison of pulmonary and extrapulmonary tuberculosis in Nepal-a hospital based retrospective study. *BioMed central infectious disease* 2008;
2. Acharya D, Majra JP. A clinical-epidemiological study of tuberculosis among hospitalized cases in Dakshina Kannada district of Karnataka. *NTI Bulletin* 2007; 43/3 & 4: 43-46.
3. American thoracic society. Diagnostic standards and classification of tuberculosis In adult and children. *Am J Respir crit care med* 2000; 161: 1376-1395.
4. Sharma SK, Lawaniya S, Lal H, Singh UB, Sinha PK, DOTS center at a tertiary care teaching hospital:Lessons Learned and future directions. *Indian J Chest dis Allied Sci* 2004; 46:251-256.
5. Baburao PD, Bhaskar PS, Deepak PV. Study of tuberculosis cases under RNTCP attending designated microscopy centre at Pravara rural hospital. *Loni, Pravara Med* 2009; 4(4).
6. Kumar GS, Kumar NHN, Ramakrishna R, Jayarama S, Kotian MS. Trend of tuberculosis cases under DOTS strategy in dakshina kannada district of Karnataka, India: issue and challenges. *Iranian J Publ Health* 2009; 38(4):72-76.