



AMERICAN JOURNAL OF PHARMTECH RESEARCH

Journal home page: <http://www.ajptr.com/>

Synthesis and Antimicrobial Activity of Metal Chelates of 2-(8-Quinololinol-5-Yl)-Methyl Amino-5 -Phenyl-1, 3, 4-Thiadiazole Derivatives

Divyesh K Patel^{1*}, Arun Singh²

*1.Department of Chemistry, Govt. Geetanjali Girls P. G. Collage, Bhopal, M.P, India.

2.Rajya Shiksha Kendra Pustak Bhawan, Arera Hills, Bhopal, M.P., India

ABSTRACT

The novel metal chelates of 2-(8-quinolinol-5-yl)-methyl amino-5 -phenyl-1, 3, 4-thiadiazole (**1**) has been synthesized and their antimicrobial properties were evaluated. These compounds were synthesized by reaction of 2-(8-quinolinol-5-yl)-methyl amino-5 -phenyl-1, 3, 4-thiadiazole (**1**) with metal acetate and characterized using IR, ¹H-NMR and elemental analysis. The synthesized compounds were screened for their *in vitro* antimicrobial activity against microorganism *P. Expansum*, *B. Thiobromine*, *Nigras pora Sp.*, *T. roseum*, *A. Niger*, *B. megaterium*, *S. aureus*, *P. aeruginosa* and *E. coli*. All of the compounds are active against the microorganism in which some are exhibited moderate to good activity, whereas some were less active.

Keywords: Metal chelates, 5-chloromethyl-8-quinolinol, 5-phenyl-1, 3, 4-thiadiazol-2-ylamine, antifungal activity, antibacterial activity

*Corresponding Author Email: divyesh.msc@gmail.com

Received 07 May 2013, Accepted 25 May 2013

Please cite this article in press as: Patel DK. *et al.*, Synthesis and Antimicrobial Activity of Metal Chelates of 2-(8-Quinololinol-5-Yl)-Methyl Amino-5 -Phenyl-1, 3, 4-Thiadiazole Derivatives. American Journal of PharmTech Research 2013.

INTRODUCTION

During recent years there has been a large investigation on different classes of thiadiazole compounds, many of which were found to possess an extensive spectrum of pharmacological activity such as antimicrobial activity¹⁻³, antimycobacterium⁴⁻⁶, anticovulsant, antitumor⁷, CNS depressants⁸, herbicidal⁹, antiviral¹⁰, antitrypanosomal activity¹¹⁻¹² and anti-inflammatory activity¹³.


8-quinolinol (8Q) and its derivatives have been introduced as chelating groups¹⁴. Some 8-hydroxyquinoline derivative has antibacterial and antifungal activity¹⁵, activity against both viral and protozoal infections¹⁶. 8Q and its metalloquinolates have attracted great interest because their high thermal stability and good electroluminescence properties make them important prototypical electron transport and emitting materials for OLED devices¹⁷⁻¹⁹. 8Q it self is inactive but exerts activity by virtue of the metal chelates produced in its reaction with metal ions in the medium. These encouraging results led us to design the chelates metal chelates of 2-(8-quinolinol-5-yl)-methyl amino-5 -phenyl-1, 3, 4-thiadiazole (**1**) which is synthesized by coupling of 5-chloromethyl-8-quinolinol with 5-phenyl-1, 3, 4-thiadiazol-2-ylamine in the presence of sodium bicarbonate followed by formation of chelates using metal acetate of different transition metal and screened for their antimicrobial activities..

MATERIALS AND METHODS

All the chemicals used were of analytical grade. 8-hydrxyquinoline, hydrochloric acid, formaldehyde, sodium bicarbonate, ethyl alcohol and different metal salt was purchased from the E. Merck (India) Limited, Mumbai. Luria broth and agar- agar were purchased from SRL, India. Acetic acid and EDTA were purchased from Sigma Chemical Co., India. The elemental contents were determined by Thermo Finigen Flash1101 EA (Italy), the metals were determined volumetrically by Vogel's method. Infrared spectra of the synthesized compounds were recorded on Nicolet760 FT-IR spectrometer. NMR spectrum of compound was recorded on 400 MHz NMR spectrophotometer. TMS was used as reference.

Synthesis of 2-(8-quinolinol-5-yl)-methyl amino-5 -phenyl-1, 3, 4-thiadiazole 1

5-chloromethyl-8-quinolinol hydrochloride (2.3 gm, 0.01 mole) and 2-amino-5-phenyl-1, 3, 4-thiadiazole (3.34 gm, 0.01 mole) were suspended in acetone (50 ml). Sodium bicarbonate (1.68 gm, 0.02 mole) was added to the reaction mass and warmed for 6 h. Solvent was evaporated and product was collected by filtration further purified by washing with acetone



Scheme-1:- synthesis of metal chelates of 2-(8-quinolinol-5-yl)-methyl amino-5-phenyl-1,3,4-thiadiazole derivatives

Synthesis of metal chelates of compound 2a-e

The metal chelates of compound with Cu^{2+} , Ni^{2+} , Co^{2+} , Mn^{2+} , Zn^{2+} and metal ions were prepared in two steps. All the metal chelates were prepared in an identical procedure. The details are given as follows.

Preparation of compound solution

Ligand 2-(8-quinolinol-5-yl)-methyl amino-5-phenyl-1,3,4-thiadiazole (0.05 mol) was taken in 500 mL beaker and formic acid (85% v/v) was added up to slurry formation. To this slurry, water was added till the complete dissolution of compound. It was diluted to 100 mL.

Synthesis of metal-chelates of 2-(8-quinolinol-5-yl)-methyl amino-5-phenyl-1,3,4-thiadiazole

In a solution of metal acetate (0.005 mol) in water (100 mL), 20 ml of above mentioned compound solution (*i.e.* containing 0.01 M compound) was added with vigorous stirring at room temperature. The pH was adjusted around 4.5 to 6 for complete precipitation of metal chelate. The precipitates were digested on a boiling water bath. The precipitates of chelate were filtered off, washed by 1:1 mixture of water: ethanol and finally with acetone and dried at 70°C for 24 hours.

Table. 1:- Physical parameter of synthesized compound 2a-e

Sr. No	Chelates	Molecular formula	M. Wt	Yield %	Colour
1	2a	$\text{C}_{36}\text{H}_{26}\text{N}_8\text{O}_2\text{S}_2\text{Cu}^{+2}\cdot 2\text{H}_2\text{O}$	765	72	Greenish Blue
2	2b	$\text{C}_{36}\text{H}_{26}\text{N}_8\text{O}_2\text{S}_2\text{Ni}^{+2}\cdot 2\text{H}_2\text{O}$	761	74	Greenish
3	2c	$\text{C}_{36}\text{H}_{26}\text{N}_8\text{O}_2\text{S}_2\text{Mn}^{+2}\cdot 2\text{H}_2\text{O}$	757	67	Brown
4	2d	$\text{C}_{36}\text{H}_{26}\text{N}_8\text{O}_2\text{S}_2\text{Co}^{+2}\cdot 2\text{H}_2\text{O}$	761	70	Blackish
5	2e	$\text{C}_{36}\text{H}_{26}\text{N}_8\text{O}_2\text{S}_2\text{Zn}^{+2}\cdot 2\text{H}_2\text{O}$	767	76	Yellow

Antifungal activity:

The fungicidal activity of all the compounds was studied at 1000 ppm concentration *in vitro*. Plant pathogenic organisms used were *Penicillium expansum*, *Botrydepladia thiobromine*, *Nigrospora Ssp*, *Trichothecium roseum* and *Aspergillus niger*. The antifungal activity of all the compounds was measured on each of these plant pathogenic strains on a potato dextrose agar (PDA) medium such a PDA medium contained potato 200 gm., dextrose 20 gm., agar 20 gm., and water 1 liter. Five days old cultures were employed. The compounds to be tested were suspended (1000 ppm) in a PDA medium and autoclaved at 120 °C for 15 min and at 15 atm. pressure. These media were poured into sterile Petri plates and the organisms were inoculated after cooling the Petri plates. The percentage inhibition for fungi was calculated after five days using the formula given below.

$$\text{Percentage of inhibition} = \frac{100 (X - Y)}{X}$$

Where

X = Area of colony in control plate.

Y = Area of colony in test plate.

The fungicidal activity displayed by various compounds is shown in Table: 1

Antibacterial Activity:

The study has been conducted according to the method adopted by Cruickshank et al. Nutrient agar broth was melted in a water bath and cooked to 45 °C with gentle shaking to bring about uniform cooling. It was inoculated with 0.5-0.6 ml of 24 hour old culture especially and mixed well by gentle shaking before pouring on the sterilized Petri dish (25 ml each). The poured material was allowed to set (1.5 hour) and there after the “cups” was made by punching into the agar surface with a sterile cork borer and soaping out the punched part of agar. Into this “cups” 0.1 ml of test solution (prepared by dissolving 100 ml of sample in 10 ml DMF) was added by sterile micropipette. The plates were noted. The antibacterial stain use are *Bacillus megaterium*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Escherichia coli* activities of all compounds are shown in Table: 3

RESULTS AND DISCUSSION:

The toxic effect of all the complexes on fungi is shown in Table 1. The results give the following conclusions. All the complexes are toxic more or less to fungi. The substitution of phenyl rings does not have more effect on the fungicidal activity of complexes. In each series the Cu-

complexes have much toxicity. This is expected because the copper salts are mostly used as fungicides. Most of the complexes inhibit the growth of the above organisms which cause decrease in many plants. Cu⁺² metal complexes are more toxic than others and the order for is Cu⁺² > Zn⁺² > Co⁺² > Ni⁺² > Mn⁺².

The IR spectrum of the compound **1** comprises the important bands due to 8-quinolinol. The bands were observed at 1624, 1578, 1507, and 1470 cm⁻¹. The broad band due to -OH group appeared at 3800 cm⁻¹. In this band the inflections are observed at 2850 and 2940 and cm⁻¹. While the latter two might be attributed to asymmetric and symmetric vibration of CH₂ of CMQ. The NMR spectrum of compound in DMSO-d₆ indicates that the singlet of 2H at 3.72 ppm of N-CH₂-Ar group. While the singlet at 5.74 ppm due to -OH group. The aromatic protons are appeared in multiplicity at 6.88 to 7.52 ppm. The melting point is 147°C. Thus the structure of compound is confirmed as shown in Scheme 1.

The metal and C, H, N contents of metal chelates of compound are also consistent with the predicted structure. The results show that the metal: ligand (M: L) ratio for all divalent metal chelate is 1:2.

The diffuse electronic spectrum of Cu²⁺ chelates shows two broad bands at 15656 and 23372 cm⁻¹. The first band shows structures suggesting a distorted octahedral structure for the Cu²⁺ metal chelates. The higher value of the magnetic moment of the Cu²⁺ chelate supports the same. The Co²⁺ metal chelate gives rise to two absorption bands at 24655 and 19980 cm⁻¹, these absorption bands indicate an octahedral configuration of the Co²⁺ metal chelate. The spectrum of Mn²⁺ polymeric chelate comprised two bands at 18658 cm⁻¹ and 24042 cm⁻¹. The spectrum of the metal chelate of Ni²⁺ shows two distinct bands at 24122 and 15191 cm⁻¹ suggested the octahedral environment for Ni²⁺ ion.

Table: 2 Antifungal activity of metal chelates of 2-(8-quinolinol-5-yl) –methyl amino-5 - phenyl-1, 3, 4-thiadiazole

Compound code	Metal Ion	Zone of inhibition in mm at 1000 ppm (%)				
		<i>P. Expansum</i>	<i>B. Thiobromine</i>	<i>N. pora.</i>	<i>T. roseum</i>	<i>A. Niger</i>
1	-	58	66	68	65	60
2a	Cu ⁺²	77	90	80	85	84
2b	Mn ⁺²	57	58	56	64	54
2c	Co ⁺²	66	67	70	74	08
2d	Zn ⁺²	70	70	74	80	63
2e	Ni ⁺²	60	62	63	75	09

Spectral data of 2-(8-quinolinol-5-yl)–methyl amino-5 -phenyl-1, 3, 4-thiadiazole 1

Yield: 87%; colour; yellow solid; m.p. 147 °C; M.F. : C₁₈H₁₃N₄OS; M.W. 334; IR (KBr, cm⁻¹):

3400 (N-H), 2940 (C-H), 2850, 1597, 1507, 1450 (C-H Ar); NMR (400 MHz, DMSO-d₆): ppm 6.88-7.58 (m, 10H, Ar-CH), 5.74 (s, 1H, -OH), 3.72 (s, 2H, -CH₂); Elemental Analysis (C% ,H% ,N% ,S%), Calculated: 64.67, 4.19,16.76, 9.58, Found : 64.5 ,4.20,16.6,9.4

Table: 3 Antibacterial activity of metal chelates of 2-(8-quinolinol-5-yl) –methyl amino-5 - phenyl-1, 3, 4-thiadiazole

Compound code	Metal Ion	Zone of inhibition (in mm)			
		Gram + Ve		Gram -Ve	
		<i>B. megaterium</i>	<i>S.aureus</i>	<i>P. Aeruginosa</i>	<i>E. Coli</i>
1	-	20	23	15	22
2a	(L-1) ₂ Cu ⁺²	21	22	20	25
2b	(L-1) ₂ Mn ⁺²	13	15	13	12
2c	(L-1) ₂ Co ⁺²	21	14	15	16
2d	(L-1) ₂ Zn ⁺²	20	18	17	21
2e	(L-1) ₂ Ni ⁺²	17	16	14	15

CONCLUSION:

The complexes were obtained as colored powdered materials and were characterized using IR spectra, electronic spectra, and magnetic measurements. The elemental analyses were in good agreement with the complexes. From the antimicrobial activity data, it is observed that the complexes exhibit higher activity than the free ligands and the metal salt. The increase in antimicrobial activity of the complexes may be due to the metal chelation. From comparative analysis as shown in Table 1 & 2 it is observed that all the metal complexes are more potent biocidal than the ligand. From the results it is clear that Cu²⁺ is highly active among the complexes of the respective metal also Cu²⁺ is most active among all which may be due to combine effect of Cu²⁺ and functional groups on the ligand.

REFERENCES:

1. Srivastava AK, Khare RK, Singh BK, Singh H. Synthesis and fungicidal activity of some 2,6-diaryl-1,3,4-thiadiazolo [3,2-b]-S-triazine-5,7-dithiones. Indian J Heterocyclic chemistry 2007;17: 109-112.
2. Foroumadi A, Mansouri S, Kinai Z, Rahmani A. Synthesis and *in vitro* antibacterial evaluation of N- [5-(5-niro-2-thienyl)-1, 3, 4-thiadiazole-2-yl] piperazinyl quinolones. Eur J Med Chem 2003; 38: 851-854.
3. Ansari KF, Lal C. Synthesis and evaluation of some new benzimidazole derivatives as potent antimicrobial agents. Eur J Med Chem 2009; 44: 2294- 2299.
4. Foroumadi A, Asadipour A, Mirzaei M, Karimi J, Emami S. Synthesis, evaluation of *in Vitro* antituberculosis activity and cytotoxicity of some 2-(5- nitro-2-furyl)-1, 3, 4-

- Thiadiazole derivatives. *IL Farmaco* 2002; 57: 765-769.
5. Talath S, Gadad AK. Synthesis and antibacterial and antitubercular activities of some 7-[4-(5-amino-[1, 3, 4] thiadiazole-2-sulfonyl)-piperazin-1-yl] fluoroquinolone derivatives. *Eur J Med Chem* 2006; 41: 918- 924.
 6. Kolavi G, Hegde V, Khazi IA, Gadad P. Synthesis and evaluation of antitubercular activity of imidazo [2, 1-b] [1, 3,4] thiadiazole derivatives. *Bioorganic & Med Chem* 2006; 14: 3069-3080.
 7. Ibrahim DA. Synthesis and biological evaluation of 3, 6-disubstituted [1, 2, 4] triazolo [3,4 b] [1, 3, 4] thiadiazoles derivatives as a novel class of potent antitumor agents. *Eur J Med Chem* 2009; 44: 2776-2781.
 8. Jatav V, Mishra P, Kashwa S, Stables JP. Synthesis and CNS depressant activity of some Novel 3-[5-substituted 1, 3, 4 -thiadiazole-2-yl]-2-styryl quinazoline-4(3*H*)-ones. *Eur J Med Chem* 2008; 43: 135-141.
 9. Cheng GW, Xing-hai L, Yong-hong L, Su-hua W, Zheng-ming L. Synthesis and Herbicidal Activity of novel sulfonyl urea containing thiadiazole moiety. *Chem Res Chinese Universities* 2008; 24: 32-35.
 10. Kritsanida M, Mouroutsou A, Marakos P, Pouli N, Garoufalias SP, Pannecouque C, Witvrouw M, Clercq ED. Synthesis and antiviral activity evaluation of some new 6-substituted 3-(1adamantyl)-1,2,4-triazolo[3,4-b] [1,3,4] thiadiazoles. *IL Farmaco* 2002; 57: 253-257.
 11. Amir M, Kumar H, Javed SA. Condensed bridgehead nitrogen heterocyclic System. Synthesis and pharmacological activity of 1, 2, 4-triazolo-[3, 4-b]-1, 3, 4- thiadiazole Derivatives of ibuprofen and biphenyl-1, 4-yloxy acetic acid. *Eur J Med Chem* 2008; 43: 2056- 2066.
 12. [12] Carvalho SA, Dasilva EF, Santa-Rita RM, De Castro SL, Fraga AM. Synthesis and Anti trypanosomal profile of new functionalized 1, 3, 4-thiadiazole-2- arylhydrazone derivatives, designed as non-mutagenic meglumine analogues. *Bioorganic & Med Chem Letters* 2004; 14: 5967-5970.
 13. Palaska E, Sahin G, Kelicen P, Durlu NT, Altinok G. Synthesis and anti-inflammatory activity of 1-acylthiosemicarbazides 1,3,4-oxadiazoles, 1,3,4- thiadiazoles and 1,2,4-triazole-3- thiones. *IL Farmaco* 2002; 57: 101-107.
 14. R. Purohit and S. Devi, Separation and determination of trace amounts of zinc and cadmium by on-line enrichment in flow injection flame atomic absorption spectrometry.

- Analyst 1991; 116: 825-830.
15. J. Giménez-Izquierdo, J. Guiteras, A. Izquierdo and M. D. Prat., Fres. Spectrofluorimetric determination of clioquinol in pharmaceutical preparations. *J. Anal. Chem.* 1991;341, 638.
 16. W. Rohde, P. Mikelens, J. Jackson, J. Blackman, J. Whitcher and W. Levinson. Hydroxyquinolines inhibit ribonucleic acid-dependent deoxyribonucleic acid polymerase and inactivate Rous sarcoma virus and herpes simplex virus. *Antmicrob. Agents Chemother.* 1976; 10(2), 234.-240
 17. Z. L. Shen, P. E. Burrows, V. Bulovic, S. R. Forrest and M. E. Thompson, Three-Color, Tunable, Organic Light-Emitting Devices *Science.* 1997; 276(5321), 2009-2011.
 18. H. Aziz, Z. D. Popovic, N. X. Hu, A. M. Hor and G. Xu, Degradation Mechanism of Small Molecule-Based Organic Light-Emitting Devices *Science.* 1999; 283(5409), 1900-1902.
 19. S. Barth, P. Muller, H. Riel, P. F. Seidler, W. Rieß and H. Vestweber Electron mobility in tris(8-hydroxy-quinoline)aluminum thin films determined via transient electroluminescence from single- and multilayer organic light-emitting diodes *Journal of Applied Physics.* 2001; 89, 3711.