



AMERICAN JOURNAL OF PHARMTECH RESEARCH

Journal home page: <http://www.ajptr.com/>

Expression, Production and Purification of Proteinases from *Alternaria* sp.

Rajamurugan. J¹, V. Muthuviveganandavel^{2*}

1. Department of plant science, Mahatma Gandhi Govt. Arts College, Mahe – 673 311, India.

2. Department of Zoology, Tagore Arts College, Puducherry-605 000, India.

ABSTRACT

Screening and expression of protease producing 5 strains of *Alternaria* sp. were obtained from the various places in Chennai, Vellore, Tamilnadu, India. The isolates were positive on tyrosin caesin nitrate agar (1%) and thus are selected as protease producing strain. The microbial growth is revealed by the mycelial dry weight determination. Maximum growth is observed in the case of *Alternaria tenuis* (0.47 mg). Equally, the maximum growth is observed in *Alternaria solani* (0.45 mg). Total protein was determined using BSA as standard. Proteinase activity was estimated using casein as the substrate. Finally the enzyme protease was purified by column chromatography. The protein was characterized using SDS-PAGE. Maximum protein content is observed in the case of *Alternaria triticola* (0.902 mg) and *Alternaria triticina* (0.753 mg). Maximum proteinase content is observed in *Alternaria triticina* (0.591 mg). This results showed that microbes under study is a good producer of extra cellular protease, which can be beneficial for industries.

Keywords: Mycelial dry weight, *Alternaria*, chromatography, protein content, Proteinase.

*Corresponding Author Email: Muthuviveganandavel6@gmail.com

Received 21 December 2013, Accepted 09 January 2014

Please cite this article in press as: Muthuviveganandavel V. *et al.*, Expression, Production and Purification of Proteinases from *Alternaria* sp. American Journal of PharmTech Research 2014.

INTRODUCTION

Enzymes are delicate protein molecules necessary for life. Proteinases are one of the most important industrial enzymes found in wide variety of microorganisms. They are molecules of relatively small size, spherical structures that catalyze the peptide bond cleavage in proteins¹. These enzymes are important in a number of biological processes viz regulation of metabolism². Gene expression, pathogenicity and biological application. It also find application in leather industry, food industry, pharmaceutical and bioremediation process^{3,4}. The largest application of proteases is in laundry detergents, where they help removing protein based stains from clothing⁵ waste processing industries^{1,6} studied the purification and characterization of serine proteases from mature coconut endosperm, while characterization of asparaginyl proteinase were investigated by⁷ The present attempt was aimed to compare the changes in the production and activity of these useful enzymes proteinases from different species of *Alternaria*.

Collection and isolation of sample

About 5 strains of *Alternaria* were obtained from the culture collection centre of Botany Department of CAH college of Melvisharam, CAS Botany, University of Madras, Dept of Biotechnology CLRI and PG. Extension Centre, Vellore and other Microbiology Laboratories nearby. A few strains were isolated from the air microflora of Kancheepuram and cultured in Tyrosin caesein nitrate agar⁸

Mycelial dry weight determination

Mycelial mats formed from microbial cultures were filtered with coarse grade glass filters with water and transferred to a previously weighed filter paper and dried at 70° C overnight and cooled at room temperature in a desiccator and weighed to a constant weight correct to the mg values given in the tables are mean of triplicate values^{9,10,11}.

Preparation of Crude extract

The microorganisms along with the culture medium were triturated in a mortar with acid-washed sand. This was centrifuged and the supernatant was collected. The residue was triturated again with distilled water, centrifuged and the supernatant was mixed with the previous extract. The total liquid collected was made up to a known volume and analyzed for proteolytic activity.

Estimation of Protein

Total protein was determined by the method of¹³ using BSA as the standard. Sample containing 50-100µg protein were mixed with 5ml of alkaline copper solution (freshly prepared with 50 ml of 1% Na₂CO₃ in 0.1 N NaOH and 1 ml of 0.5% CuSO₄. 5H₂O in 1% potasium tartrate) and

kept for 10 mins at room temperature. It was then mixed with 1 ml of Folin-Ciocalteau's phenol reagent and allowed to stand for 30 mins and the blue color developed was read at 640 nm.

Determination of Proteinase enzyme

Proteinase activity was estimated according to the method of ¹² using case in as the substrate. In this method, a protein substrate is subjected to enzymatic hydrolysis. The tyrosine liberated by the hydrolysis is quantitatively estimated by measuring the absorbance at 275 nm. The amount of tyrosine liberated is directly proportional to the enzyme activity. 1ml of casein solution (1% w/v in 0.1 m Tris-HCl buffer pH 9.0) was mixed with 0.5 ml of suitably diluted enzyme solution and incubated at 45 C for 10 min. 50% TCA was then added to arrest the enzyme action. The mixture was warmed and filtered through whatman no.1 filterpaper. The filtrate was read at 275 nm. A control was run in an identical manner except that the enzyme was added after the addition of TCA

Enzyme Purification

Ammonium Sulphate Precipitation

The organism was grown for 48 hours as described previously. The cells were separated by centrifugation (10 000 rpm, 15 minutes), and the supernatant was fractionated by precipitation with ammonium sulfate between 50% and 70% of saturation. All subsequent steps were carried out at 4°C. The protein was resuspended in 0.1M Tris-HCl buffer, pH 7.8, and dialyzed against the same buffer. Further purification of dialyzed protein was done by column of Sephadex G-200, SDS-PAGE

The elution profile of protease by Sephadex G-200 column Chromatography showed, the dialyzed ammonium sulphate precipitates loaded on to a DEAE Sephadex G 200 column. Proteins eluted are collected and tested for specific protease activity. The absorption peaks at 280 nm were observed. The column purification shows greatest activity of enzyme. Eluted samples from Sephadex column were run on a 12% SDS page stained with silver. The band obtained with the molecular weight of 60,000 Da as protease activity band.

RESULTS AND DISCUSSION

In recent times, *Alternaria* proteinases are receiving much attention, because of their increasing applications in chemical, detergent and leather industries. The new field of enzyme engineering has in a short period of time made striking contribution to industry, medicine, agriculture and in pollution control. Proteinases are obtained from selected strains of *Alternaria* molds and bacteria. Figure.1 and Table 1 shows the mycelial dryweight, protein content and proteinase activity of

different species of *Alternaria* in culture filtrate. The growth is revealed by the mycelia dryweight determination. Among 5 species of *Alternaria* tested, maximum amount of dryweight is observed in *Alternaria tenuis*. Equally good amount of growth is seen in *Alternaria solani*. The protein content is maximum in *Alternaria triticola*. Good amount of protein content is also observed in *Alternaria triticina*. The extracellular proteinase activity is maximum in *Alternaria triticina*. Good amount is also noted in *Alternaria tenuissima*. Finally the enzyme proteinase was purified by column chromatography Figure. 2 shows the elution profile of protease by G 200 column chromatography. Proteins eluted all are collected and tested for protein activity. The column purification shows greatest activity of enzyme.

Table - 1. Mycelial Dry Weight, Protein content and Proteinase activity of *Alternaria* species.

Sl. No.	Name of the Genus	Mycelial dry weight (mg)	Protein content (mg)	Proteinase activity (mg)
1	<i>Alternaria solani</i>	0.45	0.118	0.512
2	<i>Alternaria tenuissima</i>	0.38	0.445	0.582
3	<i>Alternaria tenuis</i>	0.47	0.559	0.568
4	<i>Alternaria triticina</i>	0.42	0.753	0.591
5	<i>Alternaria triticola</i>	0.39	0.902	0.513




Figure. 1 Mycelial Dry Weight, Protein Content And Proteinase Activity of *Alternaria* Species




Figure. 2 Elution profile of Protease by Sephadex G –200 Column




Figure. 3 Electrophoresis of Proteinase:

Figure. 3 indicates the electrophoresis of proteinase. Eluted samples from sephadex column were run on 12% SDS polyacrylamide gel stained with silver. The band obtained with the molecular weight of 60,000 Da as the protease activity band.

Among the strains, *Alternaria tenuissima* and *Alternaria triticina* were producing proteinases and *Bacillus* is the most useful source for the production of proteinases. Proteinase is particularly suitable for industrial use, as it can be produced economically on a large scale, by controlled fermentation (Banerjee *et al.*, 1999). While concentrating the large scale of production of proteinase enzyme only 5 species of *Alternaria* have taken for this investigation. Out of which only 2 *Alternaria* sp. are turned to be very good sources for the proteinase enzyme production. These microorganisms through cultural methods have suggested for the large-scale production of proteinase enzyme in future.

ACKNOWLEDGEMENT

The authors are thankful to Dr. R. Puvanakrishnan, Deputy Director and senior scientist in the Department of Biotechnology Central Leather Research Institute, Adyar, Chennai for the timely help of the purification of proteases in his laboratory.

REFERENCES

1. Polgar, Laszlo. Mechanisms of protease action. CRC Press, Inc., Boca Raton, Florida. 1989; 43-76.
2. Rao, M.B., A.M. Tanksale, M.S. Ghatge and V.V. Deshpande. Molecular and biotechnological aspects of microbial proteases. *Microbiol Mol Biol Rev* 1998; 62: 597.
3. Anwar, A and Saleemuddin, M. Alkaline proteases. *Bioresource Technology*, June 1998; vol. 6(3): 175-183.
4. Gupta, R., Beg, Q.K and Lorenz, P. Bacterial alkaline proteases: molecular approaches and industrial applications. *Applied Microbiology and Biotechnology*, April 2002; vol. 59(1): 15-32.
5. Banerjee, U.C., Sani, R.K., Azmi, W and Soni, R. Thermostable alkaline protease from *Bacillus brevis* and its characterization as a laundry detergent additive. *Process Biochemistry*, October 1999; vol. 35(1): 213-219.
6. Panicker, L.M., Usha, R., Roy, S., Mandal, C. Purification and characterization of a serine Protease (CESP) from mature coconut endosperm. *BMC Res Notes* 2009; 9: 2-81.
7. Oliver, E.M., Skuce, P.J., McNair, C.M., Knox., D.P. Identification and characterization of an asparaginyl proteinase (legumain) from the parasitic nematode, *Heamonchus contortus*. *Parasitology* 2006; 2: 1-8.
8. Annadurai B. Studies on endopoly galacturonase in leafblight disease of onion (*Allium cepa* Linn) caused by *Alternaria cepulae* (Ponnappa) and its interaction with phytohormones. Ph.D. thesis, University of Madras; 1989.
9. Annadurai, B and Motlag, D.B. Effect of various carbon sources on production of endopolygalacturonase of *Alternria cepulae*. *J Ecotoxicology & Environmental monitoring* 2000; 10: 37- 41.
10. Annadurai, B., Gopinath, D and Palani, R. Studies on the role of the cell wall degrading enzymes in leafblight disease of onion (*Allium cepa* Linn.) caused by *Alternaria cepulae*, *Bio J* 1998; 10: 173-178.

11. Annadurai, B., Karunanidhi, P and Mahalingam, S. Pectic enzymes of *Alternaria cepulae* in leafblight disease of onion. J Ecobiol 1999; 11: 299-305.
12. Kurnitz, M. Crystalline soybean Trypsin Inhibitor – II: General properties. Journal of General physiology 1947; 30(4): 291-310.
13. Lowry, O.H., N.J. Rosebrough, A.L. Farr and R.J. Randall. Protein measurement with the Folin phenol reagent. J Biol Chem 1951; 193: 265-275.

AJPTR is

- Peer-reviewed
- bimonthly
- Rapid publication

Submit your manuscript at: editor@ajptr.com

