



# AMERICAN JOURNAL OF PHARMTECH RESEARCH

Journal home page: <http://www.ajptr.com/>

## Eradication of Hydrocarbon which Disrupts the World Economy through Fungi

Merlyn Stephen<sup>1\*</sup>, A Panneerselvam<sup>1</sup>

1. PG & Research Department of Botany and Microbiology, A.V.V.M.Sri Pushpam College  
(Autonomous) Poondi-613503 Thanjavur Dt. Tamil Nadu, India.

### ABSTRACT

Oil spills are considered as one of the critical problems which cause a decline in environmental health. Currently the biological solutions have become more familiar to remove hazardous substances from the environment. Oil spillage is the accidental discharge or pouring of crude oil into the environment which involves the contamination of the environment with liquid hydrocarbons. These spills endanger public health, drinking water and natural resources and disrupt the economy. Negative socio-economic impacts include decreased tourism and the closure of recreational, fishing and shellfish areas. Boats and fishing gear may be damaged and human health can also be affected through direct contact or inhalation of the oil or by eating contaminated seafood. The objective of the Oil Spills Protocol is to strengthen national and regional preparedness and response capacity to oil spills. The Protocol also serves to foster and facilitate co-operation and mutual assistance among the nations and territories in cases of emergency in order to prevent and control major oil spill incidents. Crude Oil Production in India increased to 774 BBL/D/1K in October of 2014 from 757 BBL/D/1K in September of 2014. Crude Oil Production in India is reported by the U.S. Energy Information Administration. Soil samples were collected from petrol bunks and automobile workshops at Thanjavur and used for the isolation of fungi. The isolated fungus was identified as *Aspergillus*, *Penicillium*, *Fusarium*, *Mucor*, and *Trichoderma* Species using lacto phenol cotton blue staining method and cultural characteristics. The growth of fungal diversity was higher due to more carbon concentration in the hydrocarbon polluted sites and the dominant species present in all stations are *Aspergillus* and *penicillium* Species. The study focuses its attention on the survey of fungi from the zone of hydrocarbon polluted area and their potential ability to bring about degradation of hydrocarbons.

**Keywords:** Soil, pollution, hydrocarbon, biodegradation, fungi.

\*Corresponding Author Email: [merlyn387@gmail.com](mailto:merlyn387@gmail.com)

Received 23 June 2015, Accepted 15 August 2015

Please cite this article as: Merlyn S. *et al.*, Eradication of Hydrocarbon which Disrupts the World Economy through Fungi. American Journal of PharmTech Research 2015.

## INTRODUCTION

Hydrocarbons are causing wide spread pollution in both the aquatic and terrestrial environment. The petroleum industry is responsible for the generation of large amounts of organic residues, as well as for the pollution of soils, rivers and seas. One of the best approaches to restoring contaminated environments is to make use of the physiological potential of microorganisms able to degrade the pollutants in a bioremediation process. It is an attractive approach to cleaning up hydrocarbons because it is simple to maintain, applicable over large areas, cost-effective and leads to the complete destruction of the contaminant (Bento *et al.*, 2005)<sup>1</sup>. Oil spillage is the accidental discharge or pouring of crude oil into the environment which involves the contamination of the environment with liquid hydrocarbon. These spills endanger public health, drinking water and natural resources and disrupt the economy (Gesinde, *et al.*, 2008)<sup>2</sup>. Strategies for controlling environmental contamination by petroleum and its derivatives have been the subject of various studies over the past three decades. When a spillage occurs the first action is to remove the oily phase by mechanical or by physical-chemical means through the application of surfactants to disperse the layer of oil. Bioremediation is an alternative that has been used to eliminate or minimize the effects of pollutants by using microorganisms which have biodegradation potential (Atlas, 1995)<sup>3</sup>. In recent times, an increasing amount of microbiological research has been devoted to bioremediation of oil-contaminated sites using various microbial species numerous microorganisms are known for their ability to degrade hydrocarbons. The biodegradation capabilities of bacteria have been recognized, but fungi have been the subject of recent research.<sup>2,4,5,6,7,8</sup> due to their ability to synthesize relatively unspecific enzymes involved in cellulose and lignin degradation, which are capable of degrading high molecular weight, complex or more recalcitrant compounds, including aromatic structures. The full potential of biodegradation by filamentous fungi for bioremediation purposes has not been fully investigated. The use of filamentous fungi isolated from contaminated soil may offer advantages for several reasons. Owing to their ability to extend through the soil by hyphal elongation, fungi can access xenobiotics. In addition fungi are capable of growing under stressful environmental conditions. The biological treatment are cheaper than chemical and physical ones. The degradation of crude oil in soil matrix through microorganisms are able to transform petroleum hydrocarbons in less toxic substances Therefore, the main goal of this work was to evaluate the colony growth rates of filamentous fungi isolated from contaminated soil area of Thanjavur District, using different

petroleum hydrocarbons or derivatives as the only carbon source, with a view to selecting strains for future employment in bio augmentation schemes.

## MATERIALS AND METHOD

### Sources of soil sample

The five oil contaminated soil samples used for the isolation were from five different stations located at the environment. The five different stations were 1. Automobile shop at Orathanadu, Thanjavur. 2. Petrol bunk at Mariyamman Kovil, Thanjavur 3. Heavy Vehicle Work shop center at Karanthai, Thanjavur 4. Heavy vehicle workshop at New bus stand, Thanjavur. 5. Heavy vehicles workshop center at Thiruchirapalli Samples from each station were collected randomly from different locations just 1cm below the soil surface and transported to the laboratory in sterile white plastic bags and kept in a refrigerator (in order to keep the organisms viable and free from any contaminant) for further analysis.

### Physico-Chemical Analysis of Soil

The collected soil samples was characterized for its physico-chemical properties. The physico - chemical parameters were measured by standard methods. Physical and chemical parameters of soil such as pH, salinity, organic carbon, nitrogen, phosphorous and potassium were analyzed. Moisture content was estimated by finding the weight difference of known quantity of soil before and after drying in a hot air oven at 60°C for 6 hours. Soil samples after removing the debris were suspended in distilled water (1:2 w/v) and allowed to settle down the sand particles. The pH of the suspension was read using pH meter (Systronics, India), to find out the soil pH. Electrical conductivity, Cation exchange capacity (CEC) of the soil was determined by using 1 N Ammonium acetate solution as described by (Jackson, 1973). Organic carbon content was determined by adopting chromic acid wet digestion method as described by Walkley and Black, 1934, Available nitrogen (Subbiah and Asija, 1956) Available phosphorus by Brayl method as described by Bray and Kutz, 1945. Available potassium (Standford and English, 1949) Calcium (Jackson, 1973) Available micronutrients such as Zn, Cu and Mn (Lindsay and Norwell, 1978). Other nutrients such as magnesium, sodium and available iron were analyzed following the method of (Barnes 1959 and Muthuvel and Udayasoorian, 1999). The physico - chemical parameters of the soil samples were analyzed at Soil Testing Laboratory Tiruchirappalli, Tamil nadu, India.

### Methods of Microscopic Examination

For light microscope the optical equipment used were dissecting microscope, research microscope (10x and 15x eye pieces and 10 x to 100x objectives), equipment for microphotography, camera Lucida ocular, stage micrometer.

### Identification of the soil fungi

Fungal morphology were studied macroscopically by observing colony features (Colour and Texture) and microscopically by staining with lacto phenol cotton blue and observed under compound microscope for the arrangement of spores conidia and conidiophores .The fungi were identified with the help of standards manuals. Manual of soil fungi (Gillman, 1957)<sup>9</sup>. Hyphomycetes (Subramaniyan, 1971) A manual of penecillia (Raper and Thom 1949)<sup>14</sup> the genus *Aspergillus* (Raper and Fennell, 1965)<sup>13</sup>.

### Screening tests Mills *et al* 1978<sup>11</sup>

The isolated colonies were inoculated in mineral salt medium (MSM) This medium used to the identification of petroleum utilizing fungi (oil degraders)

Composition of mineral salt medium:

Sodium nitrate	-	0.25 g
Mgso4	-	0.13 g
Kcl2	-	0.12 g
Feso4	-	0.002 g
Agar	-	3.75 g

Add one ml of petrol to sterile MSM which was cooled at 45°C under aseptic condition Streptomycin is added to the medium to prevent bacterial growth MSM and crude oil was mixed and poured into the sterile petriplates and incubated at room temperature for one week.

### *In Vitro* Analysis on the Efficient Fungal Species for Biodegradation of Different Hydrocarbons

Cultured filamentous fungi are used for the *in vitro* analysis .Prepared mineral broth in conical flask (100ml) and added crude oil in different concentration (0.5 ml, 1 ml and 1.5 ml )then had checked the OD value of the broth Then inoculated the pure culture incubated at 28°C for 12 days on the third day OD value had been checked.

## RESULTS AND DISCUSSION

The soil samples examined for isolating fungi from petroleum hydrocarbon contaminated area. Totally 15 strains were isolated from the hydrocarbon polluted soil. They were identified as

*Aspergillus* sp, *Penicillium* sp, *Fusarium* sp, *Cladosporium* sp, *Mucor* sp, *Curvularia* sp, *Helminthosporium* sp, and *Trichoderma* Sp , Among them *Aspergillus* and *Penicillium* sp are the most commonly encountered genera of hydrocarbon degraders in oil contaminated stations, in agreement with the present results.

**Table1: List of Fungi Isolated From Five Different Stations**

Isolate Identification	Station I	Station II	Station III		Station IV	Station V
	Automobile shop at Orathanadu, Thanjavur.	Heavy vehicles workshop at Thiruchirapalli	Heavy Work centre at Karanthai, Thanjavur	Vehicle shop at	Heavy vehicle workshop at New bus stand, Thanjavur	Heavy vehicles workshop at Thiruchirapalli
<i>Aspergillus fumigates</i>	+	+	+		+	+
<i>A. flavus</i>	+	+	+		+	+
<i>A.niger</i>	+	+	+		+	+
<i>A. versicolor</i>	+	-	+		+	+
<i>A.japonicus</i>	+	-	+		+	+
<i>A.nives</i>	+	-	+		-	-
<i>A.terreus</i>	+	+	+		+	+
<i>Penicillium lanosum</i>	+	-	+		-	-
<i>P.chrysogenum</i>	+	+	+		+	+
<i>P.lanosum</i>	+	+	+		+	+
<i>P.corylophilum</i>	+	-	+		+	+
<i>Trichoderma viride</i>	+	-	-		-	-
<i>Cladosporium herbarum</i>	-	+	-		+	+
<i>Mucor racemosus</i>	-	+	-		+	-
<i>Helminthosporium solani</i>	+	+	-		-	-

(+) Present (-) Absent


**Table 2: Physico-Chemical Properties of Hydrocarbon Contaminated Soils**

SI NO	Soil parameters	Stations				
		I	II	III	IV	V
1	Texture Sandy Loam	7.86	7.56	7.43	7.48	7.91
2	Temperature(°C)	50	30	40	40	30
3	pH	8.5	8.0	7.0	7.5	7.0
4	Total nitrogen (mg/gm)	0.1123	0.106	0.103	0.105	0.098
5	Total phosphorous (mg/gm)	0.004	0.004	0.004	0.005	0.004
6	Total potassium (mg/gm)	0.118	0.121	0.124	0.117	0.125
7	Zinc (ppm)	0.90	0.79	0.82	0.96	1.02
8	Copper (ppm)	0.48	0.50	0.42	0.48	0.52
9	Iron (ppm)	4.87	4.58	4.63	4.56	4.62
10	Manganese(ppm)	2.16	2.58	24.6	27.1	28.6
11	Electrical conductivity (Ec)	0.48	0.52	0.26	0.42	0.27

**Table 3 Preliminary screening of Hydrocarbon degrading fungi by plate assay**

Hydrocarbon	Fungal isolates	Biodegradation of Hydrocarbon	Fungal growth Measurement (mm)
Crude oil (1%)	C-1	+	25
	C-2	+	19
	C-3	+	15
	C-4	+	23
	C-5	+	22
	C-6	+	23
	C-7	-	-
	C-8	+	22
	C-9	+	-
	C-10	+	21
	C-11	+	16
	C-12	+	15
	C-13	-	-

(+)-Present (-)-absent

**Figure 1: The Growth Pattern of Fungi in Oil-Polluted Mineral Broth**

Studies on the isolation of filamentous fungi in environments containing oil or its sub products found a very similar diversity of genera to that found in our study, such as *Aspergillus* and *Penicillium*, Recently, it was recorded that the genera of fungi such as *Penicillium*, *Aspergillus*, *Fusarium*, *Rhizopus* associated with petroleum hydrocarbon contaminated soil. In their studies they isolated *Penicillium* and *Aspergillus* from hydrocarbon contaminated soil and identified as hydrocarbon degrading fungi along with *Trichoderma*, and *Rhizopus species*. The similar results of

our study were also obtained by Obire *et al.*, 2009; in their studies on effect of different concentrations of crude oil on fungal populations of soil. The fungal isolates obtained in their study were mainly *Aspergillus* species, while others were *Penicillium*, *Rhizopus* and *Rhodotorula* species which were all able to utilize hydrocarbon as carbon source from contaminated soil, which were able to degrade crude oil. In the present investigation *Aspergillus* and *Penicillium* species were present in dominant numbers. Our finding coincides with the work of Elisane *et al.*, 2008 who also isolated four strains from the contaminated soil. They were identified as *Aspergillus* are the most commonly encounter hydrocarbon degraders in oil contaminated tropical soils, which are in agreement with the present work fungi from total hydrocarbon contaminated soil and identified by microscopy as *Penicillium*, *Aspergillus* The different result from our findings were obtained who also isolated many fungal species that were able to degrade polycyclic aromatic hydrocarbons. The species isolated were *Coniothyrium fuckelii*, *Gliocladium virens* *Phialophora hoffmannii*, *Scopulariopsis brumptii* *Trichoderma harzianum* along with genera were similar to our finding. The growth rate of each fungus shows that *Rhizopus sp.* had the highest growth diameter in low petro contaminated PDA media culture and *Aspergillus* and *penicillium* species had the highest growth diameter in high petro contaminated PDA medium. The potentiality of the fungal isolates to degrade hydrocarbon compounds (crude oil) were screened for 21 days and the degradation percentage of top degrader were determined as 96.5 and 94.8 per cent for crude oil respectively (Table 3). Liquid culture characteristics of all the isolates during the period of degradation were summarized in Table 4, which give an indication regarding the qualitative change in the medium during degradation by the isolates.

## CONCLUSION

Biological recovery processes of soils contaminated by hydrocarbons and their derivatives have been based on the stimulation of native microorganisms and, in some cases, on the increase of the microbial population, through incorporation of native/exogenous organisms. These microorganisms do not seem to be competitive in comparison with the native microbial population, which is already adapted to the environment. One alternative would be the isolation of species from contaminated soil and their posterior growth and reintroduction into the same system. The problem is that conventional isolation methods are only able to extract a small part of viable microorganisms from the environment, thus limiting the achievement of species of interest. This fact will lead to a future development of better studies with these fungi, as well as with those that grow in both conditions, for specific purposes of use in biodegradation. In the present study the

*Penicillium* and *Aspergillus* sp was the most frequent genus found in uncontaminated soil. However, it was not isolated from contaminated soil. The existence of species that grow with or without the presence of oil allow inferences on their use as an contamination indicator or on how these hydrocarbons are degraded, having potential for the treatment of environments and the most efficient strains were used further for eradicating hydrocarbon pollution in the environment and will help to prevent various socio economical impacts.

## ACKNOWLEDGEMENTS

The authors gratefully acknowledge the Secretary and correspondent, Principal and Dean Faculty of science for their support for providing laboratory and working facilities.

## REFERENCES

1. Bento FM, Camargo FAO, Okeke BC, Frankenberger WT, Comparative bioremediation of soils contaminated with diesel oil by natural attenuation, biostimulation and bioaugmentation. *Bioresour. Technol.* 2005; 96: 1049-1055.
2. García IG, Peña PRJ, Venceslada JLB, Martín AM, Santos MAM, Gómez ER *et al.* Removal of phenol compounds from olive mill wastewater using *Phanerochaete chrysosporium*, *Aspergillus niger*, *Aspergillus terreus* and *Geotrichum candidum*. *Process Biochem.* 2000; 35: 751-758.
3. Atlas, RM. Bioremediation of Petroleum Pollutants. *Int. Biodet. Biodegradation* 1995: 317-327.
4. Colombo JC, Cabello M, Arambarri AM. Biodegradation of aliphatic and aromatic hydrocarbons by natural soil microflora and pure cultures of imperfect fungi. *Environ. Pollut* 1996; 94: 355- 362.
5. Krivobok S, Miriouchnik E, Seigle-Murandi F, Benoit-Guyod JL. Biodegradation of anthracene by soil fungi. *Environ. Pollut* 1998; 37: 523.
6. Salicis F, Krivobok S, Jack M, Benoit-Guyod JL. Biodegradation of soil fungi. *Environ. Pollut.* 1999; 38: 3031-3039.
7. Garon D, Krivobok S, Seigle-Murandi F. *Fungal degradation* 2000; 40: 91-97.
8. Potin O, Rafin C, Veignie E, Bioremediation of an aged polycyclic aromatic hydrocarbons (PAHs) – contaminated soil by filamentous fungi isolated from the soil. *Int. Biodeterior Biodegradation* 2004; 54: 45-52.
9. Gillman, JC. *A manual of Soil Fungi Revised 2nd edition* Oxford and IBH publishing company (Indian reprint) Calcutta, Bombay, New Delhi. 1957::436.

10. Gesinde AF, E.B. Agbo M.O. Agho and E.F.C. Dike. Bioremediation of some Nigeria and Arabian crude oils by fungal isolates. *Int J Pure Applied Sci*, 2008; 2:37-44.
11. Mills AL, C. Breuil, RR. Colwell, Enumeration of petroleum – degradating marine and estuarine microorganisms by the most probable number method. *Can. J. Microbiol* 1998; 24:552-557.
12. Nwachukwu, SCU. Enhanced rehabilitation of tropical aquatic environment polluted with crude petroleum using *Candida utilis*. *J. Environ. Biol.*, 2000; **21**(3): 241-250.
13. Raper KB, Fennell DI. *The genus Aspergillus*, (The Williams and Wilkins Co., Baltimore), 19:659, 686.
14. Raper KB, Thom C. *A manual of Penicillia*. Williams and Wilkins Co., Baltimore, Md., U.S.A.

***AJPTR is***

- Peer-reviewed
- bimonthly
- Rapid publication

Submit your manuscript at: [editor@ajptr.com](mailto:editor@ajptr.com)

