



AMERICAN JOURNAL OF PHARMTECH RESEARCH

Journal home page: <http://www.ajptr.com/>

Comparative Studies On Phytochemical Screening and Metal Analysis of Hydroalcoholic Extracts of *Beta Vulgaris*, *Carica Papaya*, and *Vitisvinifera*

Chandaka Madhu¹, Dr.K.Murali Krishna¹, K.Ramanji Reddy*², Dr.Mohammed Omar³
Sunkara Bhawani¹, Sravya Kalakonda¹

1.MLR Institute of pharmacy, Dundigal, Ranga Reddy Dist, Hyderabad, TS 500043.

2 Startech Labs Pvt.Ltd, Madinaguda, Hyderabad, TS-5000050

3.Arya College of Pharmacy, Kandi, Sangareddy, Hyderabad, TS-502285

ABSTRACT

The acceptance of remedies in any system of medicine depends upon their efficacy and non-toxicity. There were many reports of rejecting Indian herbal products in various countries because of higher levels of heavy metal content than permitted. Though few are regarded as essential, some heavy metals, especially cadmium, mercury and lead, are potentially hazardous due to their intrinsic or selective toxicity, particularly in environmental contexts. In view of the importance of herbal drug standardization, it was contemplated to carry out the heavy metal determination in papaya fruits (*Carica papaya*), grapes (*Vitisvinifera*) and *Beta vulgaris* (beet root) which were procured from local market. The metals like copper, iron, magnesium nickel, sodium, potassium, lead, chromium, zinc etc were estimated in the hydroalcoholic extracts by using Atomic Absorption Spectroscopy (AAS). Measurements were made using a hollow Electron Discharge Lamp (EDL). These extracts were also subjected to general phytochemical screening. Its an alarming bell for humankind if the heavy metal content is more than permitted in common edible commodities. Further study may require understanding the factors influencing the heavy metal content in commonly used fruits and vegetables.

Key words: *Carica papaya*, *Vitisvinifera*, *Beta vulgaris*, Atomic Absorption Spectroscopy, hollow Electron Discharge Lamp

*Corresponding Author Email: ramanjibiotech@gmail.com

Received 12 May 2017, Accepted 31 May 2017

Please cite this article as: Reddy KR *et al.*, Comparative Studies On Phytochemical Screening and Metal Analysis of Hydroalcoholic Extracts of Beta Vulgaris, Carica Papaya, and Vitisvinifera. American Journal of PharmTech Research 2017.

INTRODUCTION

A metal is a word which is derived from Greek called as métallon^{[1][2]}. Metal is a material (an element, compound, or alloy) that is typically hard, opaque, shiny, and has good electrical and thermal conductivity. About 91 of the 118 elements in the periodic table are metals. Metals are mainly divided into five types based upon its nature^{[3][4][5]}. Some metals adopt both structures depending on the temperature.^[6]

S.no	Type of metals	Nature	Example
1	Base metal	Metal that oxidizes or corrodes relatively easily, and reacts variably with dilute hydrochloric acid (HCl) to form hydrogen	iron, nickel, lead and zinc
2	Ferrous metal	The term "ferrous" is derived from the Latin word meaning "containing iron".	Iron
3	Noble metal	Noble metals are metals that are resistant to corrosion or oxidation	gold, platinum, silver, rhodium, Iridium and palladium
4	Precious metal	A precious metal is a rare metallic chemical element of high economic value. Chemically, the precious metals are less reactive than most elements	Gold, silver, palladium, ruthenium, rhodium, palladium, osmium, iridium, and platinum.
5	Heavy metal	A heavy metal is any relatively dense metal or metalloid and toxic to the human body.	Arsenic, lead etc

According to IOM recommended guidelines the adequate intake of minerals are given below along with their advantageous and disadvantages [7-22]

Recommended adequate intake by the IOM for Minerals mg/day										
Age	Ca	Cr	Cu	Fe	Pb	Mg	Ni	K	Na	Zn
		mcg/day			Mcg/day					
0-6 months	200	0.2	0.20	0.27		30		400	120	2
7-12 months	260	5.5	0.22	11		75		700	370	3
1-3 years	700	11	0.7	7		80		3000-4500	1000	3-8
4-8 years	1000	15	1	10		130		4500-4700	1200	8-11
9-18 years	1300	25	1.3	8		240	100-300	4700	1500	13
19-50 years	1000	35	1.5	11		410	400-600		1500	
51-70(male)	1000	35	1.7	8		400	500-700		1500	
51-70(female)	1200	30	1.7	8		420			1200-1300	
71+ years	1200		1.7							

S.No	Name of the metal	deficiency	Toxicity
1	Ca	Insomnia, anxiety, nervousness, depression, fatigue, muscle / joint pains, muscle spasms / cramps, stomach acid, osteoporosis, seizures, birth defects, miscarriage, high blood pressure, irregular heartbeat,	Arteriosclerosis, cardiovascular disease, arrhythmia, ischemic heart disease and stroke, hypertension, low stomach acid, depression, fatigue, glaucoma, higher risk for several cancers, muscle / joint pains, osteoporosis, osteoarthritis, calcification, dry skin, constipation
2	Cr	Reduced glucose tolerance / impaired glucose metabolism, weakened immune system, increased susceptibility for infections (e.g. bladder, left tonsil), trabecular bone loss,	Spinal / joint degeneration, depressed immune system, lymphatic swelling
3	Cu	Anemia, increased susceptibility for infections, weakened immune system, hormonal disorders, increased risk for (colon) cancer, miscarriage, trabecular bone loss, inflammatory joint disease	Wilson's disease, anemia, nausea, vomiting, abdominal pain, moodiness, violent behavior, ADD / ADHD, depression, confusion, weight gain, hemangiomas, arthritis, joint / spinal degeneration,
4	Fe	Fatigue, anemia, depression, dizziness, asthma, gastrointestinal disorders, pale skin, miscarriage, amenorrhea (failure to menstruate), dysmenorrhea, (painful periods), migraine headaches, Ménière's disease,	Hemochromatosis, migraine headaches, arthritis, high blood pressure, heart disease, liver disease, dizziness, gastrointestinal disorders, nausea, higher risk for several cancers, fibroid tumors, benign prostatic hypertrophy (BPH),
5	Pb	---	lead toxicity is the nervous system, both in adults and children. damage to the brain and kidneys. reduce fertility, delayed puberty
6	Mg	Irregular heartbeat, cardiovascular disease, anxiety, insomnia, nervousness, fatigue, muscle / joint pains, osteoporosis, seizures, high stomach acid, asthma, high blood pressure,	Cardiovascular disease, arrhythmia, cardiac arrest, coma, muscle spasms, joint / spinal degeneration, bone loss, low stomach acid, low body temperature, low blood pressure, higher risk for several cancers,.
7	Ni	Hyperglycemia (high blood sugar), low blood pressure, depression, liver disease, anemia, low stomach acid, sinus congestion, fatigue, low adrenals	Angina, skin rash, hypoglycemia, decreased estrogen, shortness of breath, asthma, nausea, lowered pulse, vomiting, diarrhea, headache, stomach irritation, increased protein in urine, increased red blood cells, heart failure
8	K	Irregular and/or rapid heartbeat, palpitations, high blood pressure (hypertension), shortness of breath, asthma, heart disease, chest pains, stroke, paralysis, muscle spasms / weakness, bladder weakness, edema (water	Irregular or slow heart beat, low blood pressure, kidney disease, cystitis - bladder infections or burning, higher risk of several types of cancer, infrequent menstrual cycles, muscle spasms or cramps, ovarian cysts (right), joint / back

		retention),kidney disease,	pains,weakened immune system, impotence,
9	Na	Fatigue, depression, mental apathy, low bloodpressure, headaches, dehydration, confusion,dizziness, arthritis, kidney stones, seizures,In some cases: greater risk for LDL-related heart disease, high blood pressure, or edema.	anxiety,insomnia, irritability, reactive hypoglycemia, coma. Edema, hypertension, stroke, dizziness, gout,headaches, kidney damage, kidney stones, stomach problems, nausea, vomiting, coma
10	Zn	Decreased growth, loss of taste and smell, sterility, low sperm count, decreased wound healing, skin rash, hair loss, heart disease, liver disease, kidney disease, muscle weakness, enlarged prostate (BPH), several types of cancer, calcium spurs, paralysis, high blood pressure, arthritis,	Nausea, vomiting, dehydration, stomach ulcers, gastrointestinal problems, prostatitis, higher risk of several types of cancer, loss of libido, impotence, joint / back pain, muscle weakness / cramps, anemia, dysmenorrhea (menstrual pain), ovarian cysts (left), numbness, tingling, tremors, seizures, insomnia, irritability, weakened immune system, hair loss.

World health organization currently encourages, recommends and promotes traditional/herbal preparation in National Health Programmers because such drugs are easily available at low cost, are comparatively safe and the people have faith in such remedies. Some traditional medicine/herbal preparation with ancient formulas have been found to contain some metals, such as lead, cadmium, mercury, arsenic, lithium etc. Even though the herbal preparations are safe, but some of the herbal preparations cause serious poisoning and toxic effect, due to the preparations containing dangerous toxic drugs or metals . Both medical professional and the general public should be alerted to the potential toxicity of herbal preparation. There should be frequent monitoring of herbal preparations, containing toxic drugs or metals.

The objective of the study was to determine the concentration of metals in plants that are used in medicine by the local community. Analysis of the metal in selected plant samples was performed by Atomic Absorption Spectrophotometer (AAS). Measurements were made using a hollow electron discharge lamp (EDL) for heavy metals at different wavelengths respectively

MATERIALS AND METHOD

COLLECTION AND AUTHENTICATION OF PLANT MATERIAL

The plant materials *Carica Papaya*, *Vitis Vinifera*, and *Beta Vulgaris* were collected in the month of December 2016 from local market in Gandimaisamma, Hyderabad.

PREPARATION OF POWDER:

The plant material of *Carica Papaya*, *Vitis Vinifera*, and *Beta Vulgaris* were shade dried and then powdered with a mechanical grinder to form a coarse powder. The powder was passed through sieve no 40 and was stored in an air tight container until further use. The powder was used for the extraction process

Preparation of hydroalcoholic extract:

The hydroalcoholic extract of the plant was prepared using Maceration process. The coarse powder of plant (500g) was taken in a beaker with the water (70%) and ethanol (30%) quantity of 1000ml and was Macerated for 72hrs. During the Maceration occasional stirring were carried out. After 72 hrs, the suspension was filtered through a fine muslin cloth. The solvent was removed by heating until residues was obtained and then calculate the percentage yield.

Phytochemical Screening

S.no	Phytochemical	Name of the test
1.	Glycosides	Legals test
2	Saponins	Froth test
3	Alkaloids	Dragendorffs test
4	Tannins	Chromic test

5	Flavonoids	Shinoda test
6	Mucilage	Ruthenium red test
7	Carbohydrates	Molisch test
8	Proteins	Xanthoproteic test
9	Phytosterols	Salkowski test

Preparation of extract sample:

1g of the extract was weighed and transferred into 50 ml of beaker. Then add 5ml of conc. HNO₃ and placed on hot plate until the organic fumes were completely stopped. Then add 25 ml of water for acid digestion on hot plate. Digestion to be taken until 50% of the sample was too evaporated and remaining sample was filtered and makeup to 25 ml, and gone for furthered dilutions.

Preparation of standard:

The standard reagent 1000ppm was purchased from The National Institute of Standards and Technology (NIST) from Pune. From the standard reagent bottles (1000ppm), they were furthered diluted to 0.5, 1, 1.5 ppm.

Select the position of Hallow Discharged Lamp which are placed in the socket (capacity of the socket is max 6 lamps only) .Blank, Standards(0.5,1,1.5ppm),Blank and Sample were aspirate into AAS.

Calculation:

The amount metals(ppm)=conc of the test X dilution factor / weight of the test taken

RESULTS AND DISCUSSION

Percentage yield of the extract:

S.No	Name of The Plant	Percentage Yield (%)
1	<i>Carica papaya</i>	3.6%
2	<i>Vitis vinifera</i>	4.6%
3	<i>Beta vulgaris</i>	3.4%

Phytochemical Screening:

Name of the plant	Alk	Carb	Gly	Tan	Phytos	Flav	sapo	Pro	muci
<i>Carica papaya</i>	+	+	+	+	+	--	--	+	--
<i>Vitis vinifera</i>	+	+	+	+	+	+	+	+	--
<i>Beta vulgaris</i>	+	+	+	+	+	+	--	--	+

ESTIMATION OF METALS:

Lamp position setup: 1

Socket	Element	Lamp type	Lamp id	Judge	Life time	Used time	Units
1	Zn	Normal	Zn-1	Ok	5000	266.5	mA*hrs
2	Cu	Normal	Cu-1	Ok	5000	277.0	mA*hrs
3	Ni	Normal	Ni-1	Ok	5000	166.4	mA*hrs
4	Cr	Normal	Cr-1	Ok	5000	112.7	mA*hrs
5	Pb	Normal	Pb-1	Ok	5000	303.3	mA*hrs

6	Ca	Normal	Ca-1	Ok	5000	382.6	mA*hrs
7	Na	Normal	Na-1	Ok	5000	516.1	mA*hrs
8	Mg	Normal	Mg-1	Ok	5000	451.5	mA*hrs
9	K	Normal	K-1	Ok	5000	340.0	mA*hrs
10	Fe	Normal	Fe-1	Ok	5000	719.5	mA*hrs

Weight of the extract taken for acid digestion:


S.no	Name of the extract	Weight taken
1	<i>Carica papaya</i>	1.069 g
2	<i>Vitis vinifera</i>	1.054 g
3	<i>Beta vulgaris</i>	1.0512 g

PARAMETERS OF METAL ANALYSIS IN AAS-6300 MODEL


Parameters of AAS for each metals										
Element	Zn	Ni	Fe	K	Mg	Na	Ca	Pb	Cr	Cu
Wavelength	213.9	232.0	248.3	766.5	285.2	589.0	422.7	283.3	357.9	324.8
Peak (nm)	213.87	232.47	248.43	766.55	285.07	588.76	422.60	283.23	358.00	324.67
Lamp current low(ma)	8	12	12	10	8	12	10	10	10	6
Slit width (nm)	0.7	0.2	0.2	0.7	0.7	0.2	0.7	0.7	0.7	0.7
Lamp mode	BGC- D2	BGC- D2	BGC- D2	NON- BGC	BGC- D2	NON- BGC	BGC- D2	BGC- D2	BGC- D2	BGC- D2
Line search	Ok	Ok	Ok	Ok	Ok	Ok	Ok	Ok	Ok	Ok
Beam balance	Ok	Ok	Ok	Ok	Ok	Ok	Ok	Ok	Ok	Ok

Selection of hollow cathode lamps and its calibration graphs for each metal:


1. Estimation of zinc:




2. Estimation of Nickel:




3. Estimation of Iron:




4. Estimation of Potassium:




5. Estimation of Magnesium:




6. Estimation of Sodium:



7. Estimation of Calcium:




8. Estimation of Lead:



9. Estimation of Chromium:




10. Estimation of Copper:



Metal concentration in ppm level in plant extract by using AAS

comparative study of metal analysis				
S.NO	Elements	Beta Vulgaris (PPM)	Carica Papaya (PPM)	Vitis Vinifera (PPM)
1	Zinc	20.29	18.25	7.239
2	Copper	3.629	2.82	23.70
3	Nickel	1.812	1.075	1.20
4	Chromium	0.980	1.131	1.704
5	Lead	13.40	4.080	6.33

6	Calcium	76.12	96.12	55.27
7	Sodium	45.69	42.37	3.57
8	Magnesium	8.835	9.241	10.73
9	Potassium	9.141	10.139	6.34
10	Iron	29.31	17.30	12.64



DISCUSSION AND CONCLUSION:

The hydroalcoholic extracts of *Carica papaya*, *Vitis vinifera* and *Beta vulgaris* were prepared and their percentage yields in ascending order are *Beta vulgaris*(3.4%),*Carica papaya* (3.6%) and *Vitis vinifera* (4.6%).

These extracts were subjected to preliminary phytochemical screening and *Carica papaya* extract consists of alkaloids, carbohydrates, glycosides, tannins, phytosterols, proteins, and mucilage. *Vitis vinifera* extract was found to contain alkaloids , carbohydrates , glycosides, tannins, phytosterol, flavanoids , saponins and proteins. Alkaloids, carbohydrates , glycosides, tannins, phytosterol, flavanoids , and mucilage were present in *Beta vulgaris* extract.

AAS can be employed to determine over 70 different elements in solution or directly in solid samples and has wide application in the fields of pharmacology, biophysics and toxicology research.

In our present study, we have estimated the metals like Zinc, Copper, Nickel, Chromium, Lead Calcium, Sodium, Magnesium, Potassium and Iron in the extracts of *Carica papaya*, *Vitis vinifera* and *Beta vulgaris*.

Zinc:

Zinc is an essential trace element for humans, animals, microorganisms and for plants. It is the second most abundant transition metal in organisms after iron. Most zinc is in the brain, muscle, bones, kidney, and liver, with the highest concentrations in the prostate and parts of the eye. It is the only metal which appears in all enzyme classes and found in nearly 100 specific enzymes. It is considered as of "exceptional biologic and public health importance", especially regarding prenatal and postnatal development

Semen is particularly rich in zinc, which is a key factor in prostate gland function and reproductive organ growth.

Symptoms of mild zinc deficiency include depressed growth, diarrhea, impotence and delayed sexual maturation, alopecia, impaired appetite, and reproductive teratogenesis etc. The U.S. Food and Drug Administration (FDA) has stated that zinc damages nerve receptors in the nose, which can cause anosmia.

Excess zinc can be harmful, and cause zinc toxicity to occur at ingestion of greater than 225 mg of Zinc. Excessive absorption of zinc can suppress copper and iron absorption.

The U.S. Food and Drug Administration (FDA) has stated that zinc damages nerve receptors in the nose, which can cause anosmia.

Tolerable Upper Intake Level UL: 4mg -40mg

Our experiments revealed that zinc was present at 20.29 ppm in *Beta vulgaris*, 18.25ppm in *Carica papaya* and 7.239 ppm in *Vitis vinifera* levels in these extracts.

These levels of zinc are far lower than the toxicity level.

Calcium

Calcium is essential for living organisms, in particular in cell physiology, where movement of the calcium ion into and out of the cytoplasm functions as a signal for many cellular processes. As a major material used in mineralization of bone, teeth and shells, calcium is the most abundant metal by mass in many animals.

Symptoms of hypocalcemia include numbness and tingling in the fingers, muscle cramps, convulsions, lethargy, poor appetite, and abnormal heart rhythms if left untreated, calcium deficiency leads to death.

Inadequate calcium intake causes osteopenia which if untreated can lead to osteoporosis. The risk of bone fractures also increases, especially in older individuals.

Excessive consumption of calcium carbonate antacids/dietary supplements (such as Tums) over a period of weeks or months can cause milk-alkali syndrome, with symptoms ranging from

hypercalcemia to potentially fatal renal failure. Persons consuming more than 10 grams/day of CaCO₃ (=4 g Ca) are at risk of developing milk-alkali syndrome,

Tolerable Upper Intake Level UL of calcium is: 1000mg -3000mg.

The experimental results indicated that **76.12**ppm in *Beta vulgaris*, **96.12** ppm in *Carica papaya* and **55.27**ppm in *Vitis vinifera* were present.

Therefore the levels of calcium present in these three tested samples are safe and give only beneficial effects.

Iron

Iron is a necessary trace element found in nearly all living organisms. Iron-containing enzymes and proteins, often containing heme prosthetic groups, participate in many biological oxidations and in transport.

The World Health Organization (WHO) estimates that approximately half of the 1.62 billion cases of anemia worldwide are due to iron deficiency.

Large amounts of ingested iron can cause excessive levels of iron in the blood. High blood levels of free ferrous iron react with peroxides to produce free radicals, which are highly reactive and can damage DNA, proteins, lipids, and other cellular components. Thus, iron toxicity occurs when there is free iron in the cell, which generally occurs when iron levels exceed the capacity of transferrin to bind the iron

Tolerable Upper Intake Level UL: 40mg- 45mg

The experimental results indicated that **29.31**ppm in *Beta vulgaris*, **17.30** ppm in *Carica papaya* and **12.64**ppm in *Vitis vinifera* were present.

These provide beneficial effects and do not cause any toxicity.

Copper

Numerous antimicrobial efficacy studies have been conducted in the past 10 years regarding copper's efficacy to destroy a wide range of bacteria, as well as influenza A virus, adenovirus, and fungi.

Copper is also found in many superoxide dismutases, proteins that catalyze the decomposition of superoxides, by converting it (by disproportionation) to oxygen and hydrogen peroxide:

Deficiency of copper in animals has resulted in anemia, osteoporosis, delayed wound healing and the development of aortic aneurysms, and loss of hair color.

Acute copper toxicity, such as that following the ingestion of more than 15 mg of elemental copper, has been associated with nausea, vomiting, diarrhea, and intestinal cramps. Intravascular hemolysis has occurred with larger ingestions.

Tolerable Upper Intake Level UL: 1mg- 10mg

The experimental results indicated that **3.629** ppm in *Beta vulgaris*, **2.82** ppm in *Carica papaya* and **23.70** ppm in *Vitis vinifera* were present.

The amounts are within the limits.

Chromium

Humans require chromium in trace amounts, although its mechanisms of action in the body and the amounts needed for optimal health are not well defined Chromium is known to enhance the action of insulin, a hormone critical to the metabolism and storage of carbohydrate, fat, and protein in the body

Few serious adverse effects have been linked to high intakes of chromium, so the Institute of Medicine has not established a Tolerable Upper Intake Level (UL) for this mineral . A UL is the maximum daily intake of a nutrient that is unlikely to cause adverse health effects. It is one of the values (together with the RDA and AI) that comprise the Dietary Reference Intakes (DRIs) for each nutrient

Tolerable Upper Intake Level: 200 mcg -5,000 mcg

The experimental results indicated that **0.980** ppm in *Beta vulgaris* **1.131** ppm in *Carica papaya* and **1.704** ppm in *Vitis vinifera* were present.

The quantities are in safe limits.

Nickel

Nickel plays important roles in the biology of microorganisms and plants. Plant enzyme urease (an enzyme that assists in the hydrolysis of urea) contains nickel.

Nickel deficiency causes hyperglycemia (high blood sugar), low blood pressure, depression, liver disease, anemia, low stomach acid, sinus congestion, fatigue, low adrenals

Tolerable Upper Intake Level UL: 200 mcg -1,000 mcg

Most of the nickel absorbed every day by humans is removed by the kidneys and passed out of the body through urine or is eliminated through the gastrointestinal tract without getting absorbed. Nickel is not a cumulative toxicant; however, larger doses or chronic exposure may be dangerous for human health and may represent an occupational hazard due to their acute toxicity and carcinogenicity.

Tolerable Upper Intake Level UL: 200 mcg -1,000 mcg

The experimental results indicated that **1.812** ppm in *Beta vulgaris* **1.075** ppm in *Carica papaya* and **1.20** ppm in *Vitis vinifera* were present.

The amounts of nickel are well within the UL values therefore safe to consume.

potassium

Potassium levels influence multiple physiological processes Resting cellular-membrane potential and the propagation of action potentials in neuronal, muscular, and cardiac tissue.

Potassium Deficiency causes irregular and/or rapid heartbeat, palpitations,high blood pressure (hypertension), shortnessof breath, asthma, heart disease, chest pains,stroke, paralysis, muscle spasms / weakness,bladder weakness, edema (water retention),kidney disease, liver disease, endometriosis,frequent menstrual cycles, high blood sugar,weight gain, fatigue, impotence.

Tolerable Upper Intake Level UL: 300mg -15,000mg

The experimental results indicated that **9.141** ppm in *Beta vulgaris* **10.139**ppm in *Carica papaya* and **6.34** ppm in *Vitis vinifera* were present.

Our estimation of potassium revealed that the amounts of potassium are well within the UL valus therefore safe to consume.

Sodium

In humans, sodium is an essential nutrient that regulates blood volume, blood pressure, osmotic equilibrium and pH; the minimum physiological requirement for sodium is 500 milligrams per day.[50] Sodium chloride is the principal source of sodium in the diet, and is used as seasoning

Sodium deficiency leads to fatigue, depression, mental apathy, low bloodpressure, headaches, dehydration, confusion,dizziness, arthritis, kidney stones, seizures,In some cases: greater risk for LDL-related heart disease, high blood pressure, or edema.

Tolerable Upper Intake Level UL : 1,500mg -2,300mg

Excess sodium intake leads to edema, hypertension, stroke, dizziness, gout,headaches, kidney damage, kidney stones, stomach problems, nausea, vomiting, coma

The experimental results indicated that **45.69**ppm in *Beta vulgaris***1. 42.37** ppm in *Carica papaya* and **3.57** ppm in *Vitis vinifera* were present.

The results proved that amounts of sodium are well within the UL valus therefore safe to consume.

Magnesium

Magnesium deficiency (hypomagnesemia) is common: and found in 2.5–15% of the general population. The primary cause of deficiency is decreased dietary intake: Other causes are increased renal or gastrointestinal loss, an increased intracellular shift, and proton-pump inhibitor antacid therapy

Health Risks from Excessive Magnesium Too much magnesium from food does not pose a health risk in healthy individuals because the kidneys eliminate excess amounts in the urine.

Tolerable Upper Intake Level UL : 65mg -360mg

The experimental results indicated that **8.835** ppm in *Beta vulgaris* **1. 9.241** ppm in *Carica papaya* and **10.73**ppm in *Vitis vinifera* were present.

Overdose from dietary sources alone is unlikely because excess magnesium in the blood is promptly filtered by the kidneys

Lead

Lead is a highly poisonous metal (whether inhaled or swallowed), affecting almost every organ and system in the body. The component limit of lead (1.0 µg/g) is a test benchmark for pharmaceuticals, representing the maximum daily intake an individual should have. However, even at this low level, a prolonged intake can be hazardous to human beings.

Recommended not more than: The component limit of lead (1.0 µg/g)

Our experimental results indicated that **13.40** ppm in *Beta vulgaris* **4.080**ppm in *Carica papaya* and **6.33** ppm in *Vitis vinifera* was present.

In consideration with reported lead toxicity with many herbal preparations and rejections of exports from India by many countries particularly the US, emphasis is given on lead amounts. . The maximum limit of lead in *Beta vulgaris* is more than permitted. However before drawing a conclusion from the results, it is to be assessed carefully based on the areas of cultivation, the effect of industrial pollution and utilization of large samples.

REFERENCES:

1. Henry George Liddell, Robert Scott, A Greek-English Lexicon, on Perseus Digital Library Metal, on Oxford Dictionaries www.empirestatemetals.com. (Metals and Minerals Mission 2007-2014)
2. Mortimer, Charles E. (1975). Chemistry: A Conceptual Approach (3rd ed.). New York:: D. Van Nostrad Company
3. Frank Kreith and Yogi Goswami, eds. (2004). The CRC Handbook of Mechanical Engineering, 2nd edition. Boca Raton. p. 12-2
4. Holleman, A. F.; Wiberg, E. "Inorganic Chemistry" Academic Press: San Diego, 2001.
5. Egbuna, Ogo I.; Bose, Anirban (2004). "Acute Aluminum Neurotoxicity Secondary To Treatment Of Severe Hyperphosphatemia Of Acute Renal Failure And The K/DOQI Guidelines: A Case Report And Review Of The Literature". The Internet Journal of Nephrology 2 (2).

6. Chugh, SN; Dushyant; Ram, S; Arora, B; Malhotra, KC (1991). "Incidence & outcome of aluminium phosphide poisoning in a hospital study". *The Indian journal of medical research* 94: 232–5.
7. Singh S, Singh D, Wig N, Jit I, Sharma BK; Singh; Wig; Jit; Sharma (1996). "Aluminum phosphide ingestion—a clinico-pathologic study". *J Toxicol Clin Toxicol* 34 (6): 703–6.
8. 10.Mathai, Ashu; Bhanu, Madhuritasingh (2010). "Acute aluminium phosphide poisoning: Can we predict mortality?". *Indian Journal of Anaesthesia* 54 (4): 302–7.
9. A Wahab, MS Zaheer, S Wahab, RA Khan. "Acute aluminium phosphide poisoning: an update" *Hong Kong Journal of Emergency Medicine*: 152.
10. Siwach, SB; Gupta, A (1995). "The profile of acute poisonings in Harayana-Rohtak Study". *The Journal of the Association of Physicians of India* 43 (11): 756–9.
11. Arsenic in drinking water seen as threat," *USAToday.com*, August 30, 2007.
12. Peter Ravenscroft, "Predicting the global distribution of arsenic pollution in groundwater." Paper presented at: "Arsenic -- The Geography of a Global Problem," Royal Geographic Society Arsenic Conference held at: Royal Geographic Society, London, England, August 29, 2007. This conference is part of The Cambridge Arsenic Project.
13. IARC Monograph, Volume 58". International Agency for Research on Cancer. 1993. Retrieved 2008-09-18.
14. "Safety and Health Topics | Cadmium". *Osha.gov*. Retrieved 2013-07-08.
15. "Plants Poisonous to Livestock - Cornell University Department of Animal Science". *Ansci.cornell.edu*. Retrieved 2012-04-09.
16. Tenenbein, Milton (2005). "Unit-Dose Packaging of Iron Supplements and Reduction of Iron Poisoning in Young Children". *Archives of Pediatrics & Adolescent Medicine* 159 (6): 557–60.
17. AAPCC Annual Reports, American Association of Poison Control Centers
18. Markowitz, Glen S.; Radhakrishnan, Jai; Kambham, Neeraja; Valeri, Anthony M.; Hines, William H.; D'Agati, Vivette D. (2000). "Lithium Nephrotoxicity: A Progressive Combined Glomerular and Tubulointerstitial Nephropathy". *Journal of the American Society of Nephrology* 11 (8): 1439–48.
19. Couper, J. (1837). "Sur les effets du peroxide de manganèse". *Journal de chimie médicale, de pharmacie et de toxicologie* 3: 223–225.

20. James, William D.; Berger, Timothy G.; Elston, Dirk M.; Odom, Richard B. (2006).
Andrews' diseases of the skin: clinical dermatology. Saunders Elsevier. p. 858.

AJPTR is

- Peer-reviewed
- bimonthly
- Rapid publication

Submit your manuscript at: editor@ajptr.com

