



AMERICAN JOURNAL OF PHARMTECH RESEARCH

Journal home page: <http://www.ajptr.com/>

Screening of Aqueous And Ethanolic Leaf Extracts of *Barleria Longiflora* L.f. For Anti-Inflammatory Activity

Rajaputana Lakshmi Manisha*, MeeraSumanth

Department of pharmacology, Visveswarapura Institute of Pharmaceutical Sciences,
Bangalore-560070, Karnataka, India.

ABSTRACT

This study was designed to evaluate, aqueous and ethanol leaf extracts of *Barleria longiflora* L.f. for anti-inflammatory activity using Formalin induced paw oedema in rats. Animals were divided into 6 groups of 6 rats. Groups 1 and 2 served as Formalin induced control and standard (Diclofenac 5 mg/Kg, i.p.) respectively. Groups 3 and 4 were treated, orally with aqueous extract of *Barleria longiflora* L.f. of 100 and 400 mg/kg b.w, respectively. Groups 5 and 6 were administered, orally with ethanol extract of *Barleria longiflora* L.f. of 100 and 400 mg/kg b.w, respectively. The paw volume was measured at 0, 1, 2, 3 and 4 hr after Formalin injection. The actual edema volume was calculated. The data was expressed as mean \pm S.E.M. The statistical analysis was done by means of ANOVA followed by Dunnett's post hock test. The aqueous and ethanol leaf extracts of *Barleria longiflora* L.f. exhibited significant reduction of edema in the rats. Ethanol leaf extract of *Barleria longiflora* L.f. revealed surpassing anti-inflammatory activity.

Keywords: *Barleria longiflora* L.f., inflammation, formalin, edema.

*Corresponding Author Email: rlmanisha10@gmail.com

Received 25 December 2017, Accepted 02 January 2018

Please cite this article as: Rajaputana LM *et al.*, Screening of Aqueous And Ethanolic Leaf Extracts of *Barleria Longiflora* L.f. For Anti-Inflammatory Activity. American Journal of PharmTech Research 2018.

INTRODUCTION

Pain and inflammation remains one of the world's major health problems¹. It is a common complaint in most patients suffering from disease conditions. Inflammation is a host defense mechanism to combat or overcome the invading pathogen or the foreign particles². Excessive inflammatory response has damaging effects, such as septic shock, which can lead to multiple organ dysfunction syndrome and death³. The side effects of steroidal and non steroidal anti-inflammatory drugs currently used for management of chronic inflammatory diseases may be more difficult to manage than the disease itself⁴. Novel potent anti-inflammatory drugs without considerable side effects from the natural sources are under investigation³. Despite the progress in modern medicine, it has been reported that more than 70% of the developing worlds population still depends on complementary and alternative systems of medicine, otherwise known as traditional medicine⁵. Herbal medicines derived from plant extracts are being increasingly utilized to treat a wide variety of clinical diseases, though relatively little knowledge about their mode of action is available. There is a growing interest in the pharmacological evaluation of various plants used in Indian traditional systems of medicine⁶. Thus the present investigation was carried out to evaluate the anti-inflammatory potential of aqueous and ethanol extracts of leaves of *Barleria longiflora* L.f. The plant *Barleria longiflora* L.f. have been traditionally used for treating cough, inflammations, dropsy, kidney stone problems⁷. The phyto constituents reported in this plant include anthraquinones, a pentacyclic titerpene arnidiol, campesterol, stigmasterol and β -sitosterol⁸.

MATERIALS AND METHOD

Plant material and Preparation of extracts:⁹

The leaves of *Barleria longiflora* L.f. (Balsaminaceae) were collected from Tirupati, India and were authenticated by Dr. K. Madhava Chetty, Sri Venkateshwara University, Tirupati, India. The authenticated aqueous and ethanolic extracts of leaves of *Barleria longiflora* L.f. were obtained from "Green Chem", Bangalore-560071. The percentage yield of extracts of *Barleria longiflora* L.f. obtained was Aqueous: 18% w/w, Ethanol 10% w/w.

Qualitative phytochemical investigation of aqueous and ethanolic extracts of *Barleria longiflora* L.f. [(Aq (BL), E (BL))]:^{10, 11}

The dry extracts were screened for the presence of various phytoconstituents responsible for the medicinal properties of the drug like alkaloids, glycosides, carbohydrate, tannins – phenolic compounds, proteins, amino acids, fats, oils, flavonoides, saponins and steroids etc.

Methodology:**Formalin induced paw oedema**^{12, 13}

Albino rats (Wistar strain) of either sex weighing between 180-200gm body weights were selected for the experimental study. They were divided into six groups of six animals each. Group 1 rats served as Formalin control (0.1ml of 2% v/v, s.c, hind paw), group 2 rats received standard drug Diclofenac 5 mg/Kg, b.w, i.p., group 3 and 4 rats were administered, orally with aqueous leaf extract of *Barleria longiflora* L.f. of 100 and 400 mg/kg b.w, respectively. Group 5 and 6 rats were administered, orally with ethanol leaf extract of *Barleria longiflora* L.f. of 100 and 400 mg/kg b.w, respectively. After 1 hr 0.1 ml of 2% v/v Formalin was injected into sub plantar region of right hind paw. The paw volume was measured at 0, 1, 2, 3 and 4hr after Formalin injection. The left paw volume was measured initially at 0 hr for all the groups which gives the normal paw volume. The difference between the left and right paw values gives the actual edema volume which was compared with Formalin control. The inhibition of inflammation was calculated using the formula,

$$\% \text{ inhibition} = 100 (1 - V_t/V_c),$$

Where 'Vc' represents edema volume in control and 'Vt' edema volume in group treated with test extracts.

STATISTICAL ANALYSIS:

The data was expressed as mean \pm S.E.M. The statistical analysis of results was done by means Analysis Of Variance (ANOVA) followed by Dunnett's post hoc test. The P value < 0.05 was considered statistically significant.

RESULTS AND DISCUSSION:

As seen from table 1, Alkaloids, Saponins, Flavonoids, Glycosides, Tannins, Phenolic compounds and steroids were present and carbohydrate and protein were absent.

As seen from table 2, in the rats administered with vehicle (Distilled water), the sub plantar injection of Formalin produced a local edema that increased progressively from 0.4333 ± 0.030 ml after the first hour to reach a maximum 0.9417 ± 0.056 within four hour. The [(Aq(BL), E(BL))] at doses of 100 and 400 mg/kg each from the third hour post formalin injection caused a dose dependent and significant ($P < 0.001$) reduction of edema in the rats. Diclofenac (5mg/kg) produced a significant ($P < 0.001$) decrease in edema at third and fourth hour when compared with the Formalin treated group.

Despite progress within medical research during the past decades, the treatment of many serious diseases remains problematic¹⁴. Chronic Inflammatory diseases remain one of the world's major

health problems¹⁵. Inflammation is generally considered as an essentially protective response to tissue injury caused by noxious physical, chemical or microbiological stimulus. It is a complex process involving various mediators, such as prostaglandins, leukotrienes and platelet activating factor¹⁶. Anti-inflammatory drugs are agents that reduce inflammation. It has been found that conventional synthetic NSAIDs accelerate damage and erosion of joint cartilage, advancing the osteoarthritis process. These NSAIDs are also known to cause liver and kidney damage with long term use⁵. Herbal medicines derived from plant extracts are being increasingly utilized to treat a wide variety of clinical diseases, though little knowledge about their mode of action is available⁶ because they are cheap, abundantly available and relatively less toxic¹⁷. Hence there is a need for the study on herbal drugs for treating inflammation for which the present study was carried out. The most widely used primary test to screen new anti-inflammatory agent measure the ability of a compound to reduce local edema induced in the rat paw by injection of formalin. The development of edema in the rat paw after the injection of formalin is a biphasic event. The initial phase of the edema is due to the release of histamine and serotonin and the edema is maintained during the plateau phase by kinin-like substance¹⁸ and the second accelerating phase of swelling due to the release of prostaglandin-like substances. Inhibition of edema observed in the formalin model may be due to the ability of extracts to inhibit these chemical mediators of inflammation¹⁹. NSAIDs were used as anti-inflammatory agents as they inhibit these different inflammatory mediators by inhibiting cyclooxygenases (COX-1 and COX-2)²⁰. prostaglandin synthesis was inhibited by Diclofenac sodium²¹. The inhibitory activity was shown by the aqueous (100,400 mg/kg) and ethanolic extracts (100,400 mg/kg) of *Barleria longiflora* L.f. The ethanolic extract has higher activity when compared to aqueous extract. The percentage inhibition for *Barleria longiflora* L.f. aqueous extract (100,400 mg/kg) was 43.29, 54.27 and etanolic extract (100,400 mg/kg) was 49.79, 58.41.

Phytochemical investigation of aqueous and ethanolic extracts of *Barleria longiflora* L.f. has revealed the presence of various phytoactive constituents such as alkaloids, saponins, flavonoids, glycosides, tannins, phenolic compounds and steroids. Previous literature studies revealed the presence of β -sitosterol²², (in *Typha angustifolia*), flavonoids³, (in *Tecoma stans*), phenolic compounds²³, (in *Polyalthea longiflora*), and tannins²⁴ (in *Ficus religiosa*) may be responsible for anti-inflammatory activity. These plant extracts also contain the similar phyto constituents which was responsible for the activity.

Phenolic compounds such as flavonoids, tannin and steroid are a major group of compounds that act as anti-inflammatory agents. Phenolics are compounds possessing one or more aromatic rings

with one or more hydroxyl groups. They are broadly distributed in the plant kingdom and are the most abundant secondary metabolites of plants, with more than 8,000 phenolic structures currently known, ranging from simple molecules such as phenolic acids to highly polymerized substances such as tannins²⁵. Phenolic compounds possess ideal structure chemistry for anti-inflammatory due to these characteristics: (1) Planar ring system which is essential in flavonoid molecules to display the activity; (2) Unsaturation in the C ring as ketonic carbonyl at C₄ and/ C₂-C₃ double bond; (3) Hydroxyl groups in B ring and at C₅ and C₇ of A ring are crucial; (4) The number and position of hydroxyl groups as the catechol group at ring B; (5) The flavones and flavonols having a hydroxyl group at 4' position of B ring showed higher activity; (6) The non-glycosylation of molecule, aglycones have a better effect than glycosides.^{26, 27} There is a relationship between phenolic structure and pro inflammatory mediators. Three main structural characters precipitate to flavonoids NO inhibitory potential:(a) a C₂-C₃ double bond, (b) a bulky group as a substituent lowered or nullify the inhibitory effect of compound (aglycones have bigger effect than glycosides), (c) 7 and 4' OH-groups, but this last feature should be accompanied by any of the aforementioned²⁸ Phenolic compounds work same as NSAIDs, additionally some of them retard other pro-inflammatory mediators besides COX by inhibiting their activity or gene expression. Besides, some phenolic compounds can up/down regulate transcriptional factors, like nuclear factor-kB (NFkB) or Nrf-2, in inflammatory and antioxidant pathways^{29, 30} Steroids exhibit the anti-inflammatory activity by regulating stability of pro-inflammatory mRNA signalling including COX-2³¹.

Table 1: Qualitative phytochemical screening of [(Aq(BL), E(BL)]

Chemical tests	<i>Barleria longiflora</i> L.f.	
	Aqueous extract	Ethanol extract
1) Test for Triterpenoids /Steroids		
Liebermann Burchard Test	+	+
2) Test for Glycosides		
Keller Killiani Test	+	+
Legals test	+	-
3) Test for Saponins		
Foam test	+	+
4) Test for Alkaloids		
Dragendorffs test	+	+
5) Test for Flavanoids		
Ferric Chloride test	+	+
Alkaline reagent test	+	+
Lead Acetate Solution test	+	+
6) Test for Tannins/phenols		
KMNO ₄ test	+	+

5% FeCl ₃	+	+
7) Test for Proteins		
Biuret test	-	-
8) Test for Free amino acids		
Ninhydrin Test	-	-
9) Test for Carbohydrates		
Molischs test	-	-
10) Test for fat and oil		
Saponification test	-	-

- Negative; + positive

Table 2: Effect of Aq(BL) AND E(BL), On formalin induced paw oedema (percentage inhibition)

Groups	0 hour		1 hour		2 hour		3 hour		4 hour	
	Paw edema (ml) Mean \pm SEM	% ROV	Paw edema (ml) Mean \pm SEM	% ROV	Paw edema (ml) Mean \pm SEM	% ROV	Paw edema (ml) Mean \pm SEM	% ROV	Paw edema (ml) Mean \pm SEM	% ROV
Control	0.4333 \pm 0.030	-	0.675 \pm 0.017	-	0.7833 \pm 0.024	-	0.8667 \pm 0.047	-	0.9417 \pm 0.045	-
Diclofenac 5 mg/kg	0.3083 \pm 0.023	-	0.395 \pm 0.063 ^{***}	41.69	0.4167 \pm 0.058 ^{***}	46.95	0.37 \pm 0.061 ^{***}	56.59	0.3117 \pm 0.042 ^{***}	68.94
Aq(BL) 100mg/kg	0.3583 \pm 0.023	-	0.5717 \pm 0.047 ^{\$}	15.84	0.575 \pm 0.070 [*]	26.34	0.5167 \pm 0.065 ^{***}	38.49	0.5167 \pm 0.065 ^{***}	43.29
Aq(BL) 400mg/kg	0.35 \pm 0.031	-	0.5333 \pm 0.030 ^{\$}	20.24	0.5383 \pm 0.028 ^{**}	30.82	0.4417 \pm 0.074 ^{***}	48.62	0.4233 \pm 0.076 ^{***}	54.27
E(BL) 100mg/kg	0.3583 \pm 0.020	-	0.5417 \pm 0.050 ^{\$}	19.90	0.5667 \pm 0.053 [*]	27.50	0.4917 \pm 0.063 ^{***}	40.82	0.445 \pm 0.073 ^{***}	49.79
E(BL) 400mg/kg	0.35 \pm 0.025	-	0.5083 \pm 0.030 ^{\$}	24.37	0.5217 \pm 0.030 ^{**}	32.74	0.405 \pm 0.046 ^{***}	52.26	0.3833 \pm 0.049 ^{***}	58.41

n=6, values are mean \pm S.E.M, one way ANOVA followed by Dunnet's post hoc test. Significance at *P<0.05,**P<0.01,***P<0.001 v/s control. Aq (BL)-Aqueous extract of leaf of *Barleria longiflora* L.f. E (BL)- Ethanollic extract of leaf of *Barleria longiflora* L.f.

CONCLUSION:

Ethanol extract of leaves of *Barleria longiflora* L.f. showed strong anti-inflammatory activity than aqueous extracts of the plants when compared with Diclofenac as standard.

ACKNOWLEDGEMENT:

The authors are thankful to VIPS Bangalore-560070, Karnataka, India, for providing technical support to this research.

REFERENCES:

1. Li RW, Myers SP, Leach DN, Lin GD, Leach G. A cross-cultural study: anti-inflammatory activity of Australian and Chinese plants. *J Ethanopharmacol* 2003; 85:25-32.
2. Benni MJ, Jayanthi MK, Suresha RN. Evaluation of the anti-inflammatory activity of *Aegle marmelos* (Bilwa) root. *Ind J Pharmacol* 2011; 43(4): 393-7.
3. Kameshwarn S, Suresh V, Arunachalam G, Frank RP, Manikandan V. Evaluation of anti-inflammatory potential of flower extract *Tecoma stans*. *Ind J Pharmacol* 2012; 44(4): 543-4.
4. Karawya MS, Ammar NM, Hifnawy MS, Al-okbi SY, Mohamed DA, El-anssary AA. Phytochemical study and evaluation of the anti-inflammatory activity of some medicinal plants growing in Egypt. *Med J Isla World Academy Sci.* 2010; 18(4): 139-150.
5. Apu AS, Bhuyan SH, Prova SS, Muhit A. Antiinflammatory activity of medicinal plants native to Bangladesh: a review. *J App Pharma sci* 2012; 2(2): 7-10.
6. Ratheesh M, Helen A. Antiinflammatory activity of *Ruta Graveolens* Linn on carrageenan induced paw edema in wistar male rats. *African J Biotech* 2007; 6(10): 1209-11.
7. Madhava CK, Shivaji K, Tulasi RK. *Flowring Plants of Chittoor District Andra Pradesh, India*, 4th edn. Students Offset Printers, Tirupati 2013; 254.
8. Rao EV, Sridhar P, Ravi JK, Vijaya TL. Anthraquinones and Arnidiol from *Barleria Longiflora* L.f. *Ind J Pharmacol* 1999; 61(5):282-6.
9. Mehta RM. *Pharmaceutics-1*. 6thed. Vallabh Prakashan, Delhi 2015; 171.
10. Satheesh KB, Suchetha KN, Vadisha SB, Sharmila KP, Mahesh PB. Preliminary phytochemical screening of various extracts of *Punica granatum* peel, whole fruit and seeds. *NUJHS* 2012; 2(4): 34-8.
11. Khandelwal KR, *Practical Pharmacognosy techniques and experiments*, 13th ed. Nirali Prakashan. Pune. 2005: 149-156.

12. Riyaz S, Manisha RL, Rajesh P, Shamshuddin M, Putta RK. Anti-inflammatory evaluation of alcoholic and aqueous extract of *Allium cepa* L in rats. Int J of Med Sci 2012; 1(6): 186-90.
13. William MC, Jeevan MBD, Nimmagadda VR, Krishna MG. Anti-inflammatory activity of methanolic extract of *Bambusa vulgaris* leaves. Int J Green Pharm 2009: 234-8.
14. Bohlin L. Structure-activity studies of natural products with anti-inflammatory effects. In :hostettmann k, editor. Phytochemistry of plants used in traditional medicine. Oxford: clarindon press; 1995.p. 137-61.
15. Yesilada E, Ustun O, Sezik E, Takaishi Y, Ono Y, Honda G. Inhibitory effects of Turkish folk remedies on inflammatory cytokines: interleukin-1 and tumor necrosis factor. J Ethnopharmacol. 1997; 58: 59-73.
16. Gryglewski RJ. Molecular mechanisms of inflammation. Eur J Rheumatol Inflamm. 1981; 4: 153-9.
17. Sarkar D, Dutta A, Das M, Sarkar k, Mandal C, Chatterjee M. Effect of *aloe vera* on nitric oxide production by macrophages during inflammation. Ind J Pharmacol. 2005; 37(6): 371-375.
18. Chauhan O, Godhwani JL, Khanna NK, Pendse VK. Anti-inflammatory activity of muktashuktibhasma. Ind J Exp Biol. 1998; 36: 985-9.
19. Carey WM, Dasi JMB, Rao NV, Gottumukkala KM. Anti-inflammatory activity of methanolic extract of *Bambusa vulgaris* leaves. Int J Green Pharm. 2009; 234-8. DOI:10.4103/0973-8258.5622.
20. Gilman AG, Hardman JG, Limbird LE. The Pharmacological Basis of Therapeutics. 10thed. New York: McGraw-Hill publishing corporation 2011: 687.
21. Tripathi KD. Essentials of Medical Pharmacology, 6th ed. Jaypee brother Medical Ltd. New delhi. 2003: 193.
22. Varpe SS, Archana RJ, Mukta PB, Parikshit RJ. Evaluation of anti-inflammatory activity of *Typha angustifolia* pollen grains extracts in experimental animals. Ind J Pharmacol. 2012; 44(6): 788-91.
23. Mandal S, Rajini GP, Sharma RK, Gupta N. In vitro antioxidant and anti inflammatory potential of *Polyalthea longiflora* in rats. Ind J Pharmacol. 2012; 44(2): 277-8.
24. Gulecha V, Sivakumar T, Upaganlawar A, Mahajan M, Upasani C. Screening of *Ficus religiosa* leaves fractions for analgesic and anti-inflammatory activities. Ind J Pharmacol. 2011; 43(6): 662-6.

25. Jin D, Russell JM. Plant phenolic: Extraction, Analysis and their antioxidant and anticancer properties. *Molecules* 2010; 15: 7313-52.
26. Gautam R, Jachak, SM. Recent developments in anti-inflammatory natural products. *Med Res Rev.*2009; 29, 767–820.
27. Lago JH, Toledo-Arruda AC, Mernak M, Barrosa KH, Martins MA, Tibério IF et al. Structure-activity association of flavonoids in lung diseases. *Molecules.* (2014); 19: 3570-95.
28. Hämäläinen M, Nieminen R, Vuorela P, Heinonen M, Moilanen E. Anti-inflammatory effects of flavonoids: Genistein, kaempferol, quercetin, and daidzein inhibit STAT-1 and NF-κB activations, whereas flavone, isorhamnetin, naringenin, and pelargonidin inhibit only NF-κB activation along with their inhibitory effect on iNOS expression and NO production in activated macrophages. *Mediators Inflamm.* 2007; 10. doi:10.1155/2007/45673.
29. Maroon JC, Bost JW, Maroon A. Natural anti-inflammatory agents for pain relief. *Surg Neurol Intern.* 2010; 1: 1–80.
30. Sergent T, Piront N, Meurice J, Toussaint O, Schneider YJ. Anti-inflammatory effects of dietary phenolic compounds in an *in vitro* model of inflamed human intestinal epithelium. *Chemico-Biol Interact.* 2010; 188: 659–67.
31. McManus R. Mechanism of steroid action and resistance in inflammation and disease. *J Endocrinol.* 2003; 178: 1-4.

AJPTR is

- Peer-reviewed
- bimonthly
- Rapid publication

Submit your manuscript at: editor@ajptr.com

